


PCT/EP 03 / 11792  
27 APR 2005



### Bescheinigung

### Certificate

### Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten internationalen Patentanmeldung überein.

The attached documents are exact copies of the international patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet international spécifiée à la page suivante.

Den Haag, den  
The Hague,  
La Haye, le

14. 11. 2003

Der Präsident des Europäischen Patentamts  
Im Auftrag  
For the President of the European Patent Office  
Le Président de l'Office européen des brevets  
p. o.

**PRIORITY  
DOCUMENT**

SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Mme. C. Rossi

Patentanmeldung Nr.  
Patent application no.  
Demande de brevet n°

PCT/EP 02/12274

BEST AVAILABLE COPY

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.:  
Demande n°:

PCT/EP 02/12274 ✓

Anmelder:  
Applicant(s):  
Demandeur(s):

1. JANSSEN PHARMACEUTICA N.V. - Beerse, Belgium
2. PEETERS, Pieter, Johan - Beerse, Belgium (US only)
3. GÖHLMANN, Hinrich, Wilhelm, Helmut - Beerse, Belgium (US only)

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

CRH RESPONSIVE GENES IN CNS

Anmeldetag:  
Date of filing:  
Date de dépôt:

31 October 2002 (31.10.2002) ✓

In Anspruch genommene Priorität(en)  
Priority(ies) claimed  
Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Benennung von Vertragsstaaten : Siehe Formblatt PCT/RO/101 (beigefügt)  
Designation of contracting states : See Form PCT/RO/101 (enclosed)  
Désignation d'états contractants : Voir Formulaire PCT/RO/101 (ci-joint)

Bemerkungen:  
Remarks:  
Remarques:

Further applicants:

4. SWAGEMAKERS, Sigrid, Maria, Alice - Beerse, Belgium (US only)
5. KASS, Stefan, Ulrich - Beerse, Belgium (US only)
6. STECKLER, Thomas, Horst, Wolfgang - Beerse, Belgium (US only)

## PCT REQUEST

PRD2009p

Original (for SUBMISSION) - printed on 31.10.2002 03:04:11 PM

IV-1	<b>Agent or common representative; or address for correspondence</b> The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as:	<b>common representative</b>
IV-1-1	Name	<b>JANSSEN PHARMACEUTICA N.V.</b>
IV-1-2	Address:	<b>Turnhoutseweg 30 B-2340 Beerse Belgium</b>
IV-1-3	Telephone No.	<b>32+14 60 21 86</b>
IV-1-4	Facsimile No.	<b>32+14 60 54 91</b>
IV-1-5	e-mail	<b>patents@janbe.jnj.com</b>
V	<b>Designation of States</b>	
V-1	<b>Regional Patent</b> (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	<b>EP: AT BE BG CH&amp;LI CY CZ DE DK EE ES FI FR GB GR IE IT LU MC NL PT SE SK TR and any other State which is a Contracting State of the European Patent Convention and of the PCT</b>
V-2	<b>National Patent</b> (other kinds of protection or treatment, if any, are specified between parentheses after the designation(s) concerned)	<b>US</b>
V-5	<b>Precautionary Designation Statement</b> In addition to the designations made under items V-1, V-2 and V-3, the applicant also makes under Rule 4.9(b) all designations which would be permitted under the PCT except any designation(s) of the State(s) indicated under item V-6 below. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit.	
V-6	<b>Exclusion(s) from precautionary designations</b>	<b>NONE</b>
VI	<b>Priority claim</b>	<b>NONE</b>
VII-1	<b>International Searching Authority Chosen</b>	<b>European Patent Office (EPO) (ISA/EP)</b>

## CRH RESPONSIVE GENES IN CNS

The present invention relates generally to therapy and diagnosis of depression. In particular this invention relates to the polypeptides as well as to the polynucleotides encoding these polypeptides, wherein said polypeptides are shown to play a central role in mediating the cellular response to corticotropin releasing hormone. These polypeptides and polynucleotides are useful in the diagnosis, treatment and/or prevention of depression.

### BACKGROUND OF THE INVENTION

Recent socioeconomic analyses found that depression is a leading cause of disability and a major risk factor for development of other diseases. Moreover, on a world-wide scale depression is underdiagnosed and undertreated. Current antidepressant drugs have proven to be effective, but are burdened with slow onset of action and side effects. Above this, it is still unclear by which pharmacological mode of action they exert their clinical effects. Hypothesis-driven research based upon the corticosteroid receptor hypothesis of depression has led to a novel concept focusing on brain neuropeptide receptors, specifically the corticotropin-releasing hormone (CRH) receptor as drug target.

Corticotropin releasing hormone (CRH), a 41-amino acid polypeptide plays a central role in the regulation of the hypothalamic-pituitary-adrenal axis, mediating the endocrine responses to various stressors. Hypothalamic neurons release CRH into the hypophyseal portal system in response to stress, stimulating the secretion and biosynthesis of pituitary adrenocorticotropin (ACTH) leading to increased adrenal glucocorticoid production (1). Several clinical and preclinical studies point towards a causal role for alterations in the CRH system in the development of depression (2). The first studies with CRH in humans showed that the ACTH response to CRH is blunted in depressed patients, reflecting a CRH receptor desensitization secondary to continuously increased hypothalamic CRH secretion (3;4). In support of blunted ACTH response as consequence of increased CRH release is the finding of elevated CRH levels in cerebrospinal fluid of patients with depression. Other findings strengthening this notion of CRH hypersecretion in the depressed state are an increased number of CRH secreting neurons and a decreased number of CRH receptors in suicide victims who suffered from depression (5;6).



-2-

For CRH two high affinity receptors have been described, CRH-R1 and CRH-R2, both of which exist in several splice variant forms. Activation of these receptors by CRH results in  $G_s$ -mediated stimulation of adenylyl cyclase leading to increased levels of intracellular cAMP. This in itself will activate cAMP dependent protein kinase A (PKA) and ultimately result in increased cytosolic levels of cAMP and  $Ca^{2+}$ . The increased levels of cAMP and  $Ca^{2+}$  lead to the activation of several other additional kinases such as  $Ca^{2+}$ /calmodulin-dependent kinase II (CAMKII) and p42/p44 mitogen activated kinases (MAPK). As a result the  $Ca^{2+}$ /cAMP response element binding protein (CREB) is phosphorylated and this in turn will regulate the transcription of genes containing cAMP response elements (CRE) in their promoter region. Examples of such genes shown to be involved in the modulation of CRH signaling include *c-fos*, macrophage migration-inhibitory factor gene *Mif*, orphan nuclear receptors *Nurr77* and *Nurr1*.

Notwithstanding the fact that the downstream pathways for CRH activated receptors were extensively studied and led to the identification of a number of genes involved in the signaling cascade, a major area is unexplored. It was thus an object of the present invention to explore the transcriptional response to CRH stimulation at a genome wide level in order to identify further genes involved in the corticotropin-releasing hormone receptor activated gene network. The polypeptides thus identified and the polynucleotides encoding said polypeptides provide new chances for drug development as drug targets through screening techniques, or are useful in the diagnosis, prevention and/or treatment of depression.

## 25 BRIEF DESCRIPTION OF THE DRAWING

Table 1 : Table of genes shown to be important mediators of CRH induced changes in the CNS.

30 Figure 1: Expression levels of CRH-R1 in different mouse tissues. Quantitative RT-PCR was applied on tissues derived from three different animals for the brain regions.

Figure 2: Spectral map analysis of microarray data obtained on all investigated brain areas. Squares depict different samples whereas circles depict genes. The diameter of the circle corresponds to the average intensity of a gene. Distances between squares are a measure for similarity between samples. A positive association of a gene with a given

sample (i.e. an upregulation of that gene in that particular sample) results in the positioning of the gene and sample on a common line through the centroid (indicated by a cross). Different areas are shown in the following order (A) cerebellum, (B) frontal cortex, (C) hippocampus, (D) nucleus accumbens, (E) temporal area, (F) pituitary.

5

Figure 3: HPA axis parameters in animals used for microarray analysis. Plasma CORT levels are 6 to 8 times higher in CRH overexpressors compared to wild-type animals. No significant differences were observed in ACTH levels.

10

Figure 4: Microarray and quantitative RT-PCR data for genes implicated in glucocorticoid signaling. (A) Array data showing a downregulation of 11 $\beta$ -hydroxysteroid dehydrogenase type 1 (HSD11b1) in hippocampus were confirmed using quantitative RT-PCR (levels normalized against  $\beta$ -actin). (B) Array data showing a significant upregulation in hippocampus and frontal cortex of FK506 binding protein 5 (Fkbp5), a modulator of glucocorticoid receptor activation. (C) Array data on serum/glucocorticoid regulated kinase (Sgk) were confirmed by quantitative RT-PCR in frontal cortex.

15

Figure 5: Microarray and quantitative RT-PCR data for neurotensin receptor 1 and 2. (A) Downregulation of neurotensin receptor 2 (Ntsr2) in CRH overexpressing mice observed in hippocampus was established using quantitative RT-PCR. Levels of neurotensin receptor 1 (Ntsr1) were below detection limit in microarray analysis. (B) Quantitative RT-PCR however established a similar downregulation of Ntsr1 in CRH overexpressors. In addition two-way ANOVA on qRT-PCR data shows a significant effect of vehicle treatment which apparently is abolished by treatment with R121919.

25

### DETAILED DESCRIPTION

As used herein, the term "compound" or "agent" means a biological or chemical compound such as a simple or complex organic molecule, a peptide, a protein or an oligonucleotide. A "test compound" as used herein, refers to a "compound" or "agent" used in a method according to the invention to assess whether said compounds modulates CRH signalling activity.

30

"CRH signaling" as used herein refers to the cellular changes in gene transcription after activation of the corticotropin releasing hormone receptor by CRH in said

35

cell. It thus induces a CRH specific gene expression profile. Changes at the transcriptional level can be assessed either at the protein level or at the gene, RNA level.

5 "CRH response activity" as used herein refers in general to the change of a detectable cellular parameter as a result of the exposure of said cell to CRH. Detectable cellular parameters include amongst others, changes in membrane potential, changes in enzyme activity of an enzyme that modulates CRH signalling a said cell, changes in expression levels of a protein according to the invention or changes in the amount of second messengers such as cGMP,  
10 cAMP,  $\text{Ca}^{2+}$  or  $\text{IP}_3$ .

The term "analog" or "functional analog" refers to a modified form of mammalian purine permeases in which at least one amino acid substitution has been made such that said analog retains substantially the same biological activity as the  
15 unmodified mammalian purine permease *in vivo* and/or *in vitro*.

"Sample" or "Biological sample" as used herein refers to cells, cell extracts, body fluids or tissue samples such as from, blood, urine, saliva, tissue biopsy or autopsy material.

The term "functional analog" is intended to include the "fragments," "variants,"  
20 "degenerate variants," "analogs" and "homologues" or to "chemical derivatives" of the polypeptides according to the invention. Useful chemical derivatives of polypeptide are well known in the art and include, for example covalent modification of reactive organic site contained within the polypeptide with a secondary chemical moiety. Well known cross-linking reagents are useful to  
25 react to amino, carboxyl, or aldehyde residues to introduce, for example an affinity tag such as biotin, a fluorescent dye, or to conjugate the polypeptide to a solid phase surface (for example to create an affinity resin)

Variant(s) of polynucleotides or polypeptides, as the term is used herein, are  
30 polynucleotides or polypeptides that differ from a reference polynucleotide or polypeptide, respectively. A variant of the polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. (1) A polynucleotide that differs in nucleotide sequence from another, reference polynucleotide. Generally,  
35 differences are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical. As noted below, changes in the nucleotide sequence of the variant may be silent. That is,

they may not alter the amino acids encoded by the polynucleotide. Where alterations are limited to silent changes of this type a variant will encode a polypeptide with the same amino acid sequence as the reference. Also as noted below, changes in the nucleotide sequence of the variant may alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Such nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed above. (2) A polypeptide that differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions, fusions and truncations, which may be present in any combination.

The terms "complementary" or "complementarity" as used herein refer to the capacity of purine and pyrimidine nucleotides to associate through hydrogen bonding to form double-stranded nucleic acid molecules. The following base pairs are related by complementarity: guanine and cytosine; adenine and thymine; and adenine and uracil. As used herein "complementary" means that the aforementioned relationship applies to substantially all base pairs comprising two single-stranded nucleic acid molecules over the entire length of said molecules. "Partially complementary" refers to the aforementioned relationship in which one of the two single-stranded nucleic acid molecules is shorter in length than the other such that a portion of one of the molecules remains single-stranded.

The term "conservative substitution" or "conservative amino acid substitution" refers to a replacement of one or more amino acid residue(s) in a parent protein without affecting the biological activity of the parent molecule based on the art recognized substitutability of certain amino acids (See e.g. M. Dayhoff, In Atlas of Protein Sequence and Structure, Vol. 5, Supp. 3, pgs 345-352, 1978).

"Fragment thereof" refers to a fragment, piece, or sub-region of a nucleic acid or protein molecule whose sequence is disclosed herein, such that said fragment comprises 5 or more amino acids, or 10 or more nucleotides that are contiguous in the parent protein or nucleic acid molecule.

"Functional fragment" as used herein, refers to an isolated sub-region, or fragment of a protein disclosed herein, or sequence of amino acids that, for example, comprises a functionally distinct region such as an active site for a receptor.

-6-

Functional fragments may be produced by cloning technology, or as the natural products of alternative splicing mechanisms.

The term "homolog" or "homologous" describes the relationship between different nucleic acid molecules or amino acid sequences in which said sequences or molecules are related by partial identity or similarity at one or more blocks or regions within said molecules or sequences. "Isolated nucleic acid compound" refers to any RNA or DNA sequence, however construed or synthesized, which is locationally distinct from its natural location.

5 A "nucleic acid probe" or "probe" as used herein is a labeled nucleic acid compound which hybridizes with another nucleic acid compound. "Nucleic acid probe" means a single stranded nucleic acid sequence that will hybridize with a single stranded target nucleic acid sequence. A nucleic acid probe may be an oligonucleotide or a nucleotide polymer. A "probe" will usually contain a detectable moiety which may be attached to the end(s) of the probe or be internal to the sequence of the probe.

10 The term "primer" is a nucleic acid fragment which functions as an initiating substrate for enzymatic or synthetic elongation of, for example, a nucleic acid molecule.

15 The term "hybridization" as used herein refers to a process in which a single-stranded nucleic acid molecule joins with a complementary strand through nucleotide base pairing.

20 The term "stringency" refers to hybridization conditions. High stringency conditions disfavor non-homologous base pairing. Low stringency conditions have the opposite effect. Stringency may be altered, for example, by temperature and salt concentration. "Stringent conditions" refers to an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 x SSC at about 65°C. Further suitable hybridization conditions are described in the examples.

25 "Lower stringency conditions" include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH<sub>2</sub>PO<sub>4</sub>; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1 X SSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC). Note that variations in the above conditions may be accomplished through the inclusion and/or

30

35

-7-

substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

The term "fusion protein" as used herein refers to protein constructs that are the result of combining multiple protein domains or linker regions for the purpose of gaining the combined functions of the domains or linker regions. This may be accomplished by molecular cloning of the nucleotide sequences encoding such domains to produce a new polynucleotide sequence that encodes the desired fusion protein. Alternatively, creation of a fusion protein may be accomplished by chemically joining two proteins.

The term "linker region" or "linker domain" or similar such descriptive terms as used herein refers to polynucleotide or polypeptide sequence that are used in the construction of a cloning vector or fusion protein. Functions of a linker region can include introduction of cloning sites into the nucleotide sequence, introduction of a flexible component or space-creating region between two protein domains, or creation of an affinity tag for specific molecule interaction. A linker region may be introduced into a fusion protein resulting from choices made during polypeptide or nucleotide sequence construction.

#### *Screening methods*

The present invention relates to screening methods to identify compounds that modulate corticotropin-releasing hormone (CRH) induced depression and stress. It is based on the identification of a number of genes as downstream modulators of the CRH activated CRH receptors. In particular this invention provides a method for identifying a compound capable to alter the CRH signalling response in a cell, said method comprising;

- a) contacting said cell with CRH in the presence and absence of said compound;
- b) determine the change at transcriptional level of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said cell; and
- c) compare the transcriptional level of said protein in the presence and absence of said compound;

whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

5

To determine the change of transcription at the protein level one could determine the amount of said protein using art known techniques. For example using separation techniques such as isoelectric focusing or SDS-page in combination with protein staining techniques such as coomassie or silver staining. Alternatively, for proteins that are enzymes, the amount in a given solution or tissue extract can be measured or assayed in terms of the catalytic effect the enzyme produces, that is the conversion of its substrate into reaction product. For example, for kinases one may assess the kinase activity using a substrate comprising the kinase specific phosphorylation site and by measuring the phosphorylation of the substrate by incorporation of radioactive phosphate into the substrate. This assay may be performed both in the presence and absence of the compound to be tested. For proteins that are not enzymes, other quantification methods are required. For example transport proteins can be assayed by their binding to the molecule they transport and hormones and toxins by the biological effect they produce.

10

15

20

To assess changes in transcription at the gene level, RNA or cDNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. Preferably said analysis method comprises the use of a labelled oligonucleotide probe targeted to a suitable region of the gene encoding a protein that modulates CRH signalling in said cell. Accordingly, in a preferred embodiment the level of gene transcription is assessed using a probe which binds to a polynucleotide encoding an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

25

30

In another embodiment, an array of oligonucleotides probes comprising a nucleotide sequence encoding a protein that modulates CRH signalling or fragments thereof can be constructed to conduct efficient screening of gene expression. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee et al., Science, Vol 274, pp 610-613 (1996)).

35

In an alternative embodiment, the method for identifying a compound capable to alter the CRH signalling response in a cell, comprises;

a) contacting said cell with CRH in the presence and absence of said compound; and

5        b) determine the expression level of a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ  
10    ID No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40.

To assess changes in expression levels, RNA or cDNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. Preferably said analysis method comprises the use of a  
15    labelled oligonucleotide probe targeted to a suitable region of the polynucleotide. Accordingly, in a preferred embodiment the level of gene transcription is assessed using a probe which binds to a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11,  
20    SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40.

In another embodiment, an array of oligonucleotides probes comprising a  
25    nucleotide sequence encoding a protein that modulates CRH signalling or fragments thereof can be constructed to conduct efficient screening of gene expression.

In case the genomic DNA is not used directly, mRNA may be isolated, and a first strand cDNA synthesis carried out. A second round of DNA synthesis can be carried out for the production of the second strand. Subsequently by the specific  
30    PCR amplification an isolated cDNA can be obtained. If desired the double-stranded cDNA can be cloned into any suitable vector, for example, a plasmid, thereby forming a cDNA library. In analogy to the above, it is possible to screen cDNA libraries constructed in a bacteriophage or plasmid shuttle vector with a labeled oligonucleotide probe targeted to any suitable region of the gene encoding a  
35    protein that modulates CRH signalling. See e.g. PCR Protocols: A Guide to Method and Application, Ed. M. Innis *et al.*, Academic Press (1990).



Methods for constructing cDNA libraries in a suitable vector such as a plasmid or phage for propagation in prokaryotic or eukaryotic cells are well known to those skilled in the art. [See e.g. Maniatis *et al. Supra*]. Suitable cloning vectors are well known and are widely available.

5 In a further embodiment changes in gene transcription are determined at mRNA level. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example; nucleic acid amplification, for instance via PCR, RT-PCR; RNase protection; Northern blotting and other hybridization  
10 methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Assay techniques that can be used to determine the presence of protein derivatives or  
15 variants comprise amongst others mass spectrometry.

It is thus an object of the present invention to provide a method for identifying a compound capable to alter the CRH signalling response in a cell, said method comprising:

- 20 a) contacting said cell with CRH in the presence and absence of said compound;  
b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said cell; and  
c) compare the amount of said protein in the presence and absence of said compound;
- 25 whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.
- 30 Preferably, the method to assay the amount of protein that modulates CRH signaling is using an antibody which binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.
- 35 Thus in another embodiment, this invention provides a monospecific antibody

immunologically reactive with a protein that modulates CRH signalling said protein being selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID  
5 No.41.

Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells expressing these to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides  
10 antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975)256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983)4:72) and the EBV-hybridoma technique (Cole *et al.*, *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, pp.77-96, Alan R.  
15 Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No.4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

20 The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against polypeptides of the present invention may also be employed to treat the CRH metabolism related disorders such as CRH induced stress or  
25 depression amongst others.

To determine the amount of protein that modulates CRH signalling, the antibodies according to the invention are used in conventional immunological techniques. Suitable immunological techniques are well known to those skilled in the art and include for example, ELISA, Western Blot analysis, competitive or sandwich  
30 immunoassays and the like, as is otherwise well known they all depend on the formation of an antigen-antibody immune complex wherein for the purpose of the assay, the antibody can be detectable labeled with, e.g. radio, enzyme or fluorescent labels or it can be immobilized on insoluble carriers.

For example in an ELISA screening format the antibody is added to a solid  
35 phase (for example the bottom of a microplate) which is coated with either the protein or a peptide fragment thereof coupled to a carrier (such as BSA), and then, adding an anti-immunoglobulin antibody (for example when the immunization is performed in

mice, an anti-mouse immunoglobulin antibody is used, e.g. sheep-anti-mouse immunoglobulin (Ig)) conjugated with a detectable label such as an enzyme, preferably horseradish peroxidase, or a radioactive isotope such as  $^{125}\text{I}$ .

It is thus an object of the invention to provide immunoassays for the  
5 determination or detection of proteins that modulate CRH signalling in a sample, the method comprising contacting the sample with an antibody to the proteins according to the invention and determining whether an immune complex is formed between the antibody and said protein. These methods can either be performed on tissue samples  
10 or body fluid samples and generally comprise obtaining a sample from the body of a subject; contacting said sample with an imaging effective amount of a detectably labeled antibody according to the invention; and detecting the label to establish the presence of proteins that modulate CRH signalling in the sample.

The measuring methods using the antibodies of the present invention are not particularly limited. Any measuring method may be used as long as the amount of  
15 antibodies, antigens or the antigens-antibody complexes corresponding to the amount of the antigens to be measured is detected by chemical or physical means, and calculated from standard curves prepared by the use of standard solutions containing the antigens in known amounts. For example, nephelometry, competitive methods, immunometric methods and sandwich methods are suitably used. With respect to sensitivity and  
20 specificity, it is particularly preferred to use sandwich methods described below.

In measuring methods using labelling substances, radioisotopes, enzymes, fluorescent substances, luminous substances, etc. are used as labelling agents. Examples of the radioisotopes include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^3\text{H}$  and  $^{14}\text{C}$ . Enzymes are usually  
25 made detectable by conjugation of an appropriate substrate that, in turn catalyzes a detectable reaction. Examples thereof include, for example, beta-galactosidase, beta-glucosidase, alkaline phosphatase, peroxidase and malate dehydrogenase, preferably horseradish peroxidase. The luminous substances include, for example, luminol, luminol derivatives, luciferin, aequorin and luciferase. Further, the avidin-biotin systems can also be used for labelling the antibodies and immunogens of the present  
30 invention.

Accordingly, in a further aspect, the present invention provides for a method of identifying and obtaining compounds that alter the CRH signalling response activity in a cell, comprising:

- 35 a) contacting a cell which expresses at least one protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ

-13-

ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41, with said test compound; and

- 5 b) compare the CRH response activity of said cell in the presence and absence of said compound.

Changes in membrane potential can be measured using conventional electrophysiological techniques and when they become available, using novel high throughput methods currently under development. Since the change in membrane potential are normally the result of ion fluxes, as an alternative approach, changes in membrane potential can be measured indirectly through the change in intracellular ion concentrations using ion-sensitive fluorescent dyes, including fluo-3, fluo-4, fluo-5N, fura red, Sodium Green, SBFI and other similar probes from suppliers including Molecular Probes. Other fluorescent dyes, from suppliers including Molecular Probes, such as DIBAC <sub>4(3)</sub> or Di-4-Anepps can detect membrane potential changes. For example calcium and sodium ion fluxes can thus be characterised in real time, using fluorometric and fluorescence imaging techniques, including fluorescence microscopy with or without laser confocal methods combined with image analysis algorithms.

20 In a preferred embodiment this assay is based around an instrument called a FLuorescence Imaging Plate Reader (FLIPR), Molecular Devices Corporation). In its most common configuration, it excites and measures fluorescence emitted by fluorescein-based dyes. It uses an argon-ion laser to produce high power excitation at 488 nm of a fluorophore, a system of optics to rapidly scan the over the bottom of a 96-/384-well plate and a sensitive, cooled CCD camera to capture the emitted fluorescence. It also contains a 96-/384-well pipetting head allowing the instrument to deliver solutions of test agents into the wells of a 96-/384-well plate. The FLIPR assay is designed to measure fluorescence signals from populations of cells before, during and after addition of compounds, in real time, from all 96-/384-wells simultaneously.

30 It is thus an object of the present invention to provide a FLIPR assay used to screen for and characterise compounds functionally active in modulating CRH response in cells, said cells expressing a protein selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

35

In an alternative embodiment, the activity of the cell may be assessed using electrophysiological methods. Therefore, proteins modulating CRH signalling in a

cell can be characterised using whole cell and single channel electrophysiology.

It is thus a further object of this invention to provide a screening method to identify compounds which modulate CRH signalling response activity in a cell, said method comprising;

- 5 (a) contacting a host cell expressing a protein selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41, with a compound to be tested;
- (b) measuring the effect of the test compound on the membrane potential of said cell
- 10 using electrophysiological techniques; and
- (c) compare the CRH response activity of said cell in the presence and absence of said compound.

In a preferred embodiment the host cells are *Xenopus* oocytes and the electrophysiological measurement consists of measuring the membrane current using

15 the voltage clamp technique at distinct membrane potentials.

Changes in enzyme activity of an enzyme that modulates CRH signalling a said cell, can generally be measured or assayed in terms of the catalytic effect the enzyme produces, that is the conversion of its substrate into reaction product. For example, for

20 kinases one may assess the kinase activity using a substrate comprising the kinase specific phosphorylation site and by measuring the phosphorylation of the substrate. Similarly for phosphatases one may assess the phosphatase activity using a phosphorylated substrate and by measuring the dephosphorylation of the substrate. These assays may be performed both in the presence and absence of the compound to be tested.

25 It is thus an object of the present invention to provide a method for identifying a compound capable to alter the CRH signalling response activity in a cell, said method comprising;

a) contacting a mixture comprising a kinase selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ

30 ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41., a source of phosphate and a suitable kinase substrate;

b) incubating said mixture in the presence or absence of a said compound and ;

measuring the level of phosphorylation of said substrate in the presence of said

35 compound compared to the level of phosphorylation of said substrate in the absence of said test compound.

In the assay of the invention, the kinase may be provided as a protein or it may be provided in the assay mixture as an mRNA encoding said kinase. When the assay comprises cell-free components, the kinase is provided as the protein. When the assay is conducted in the milieu of a cell, the kinase may be provided as either the protein or  
5 as an mRNA encoding said kinase, wherein, in order that the kinase be available in the assay, the mRNA is translated and kinase protein is thereby produced. It will be apparent from the Examples provided herein that it is a simple matter to obtain mRNA specifying the kinase and inject the mRNA into a cell for production of the kinase protein. The kinase may also be provided by expression of a plasmid, which encodes  
10 the kinase protein. Standard molecular biology techniques may be used to construct operable plasmids encoding the kinase protein and to express the plasmid in cells (Sambrook, et al., 1989, In: Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York).

As discussed herein, the method of identifying a kinase modulator may be performed  
15 either in vitro wherein the assay mixture is cell-free, in vitro wherein live cells are included in the assay, or in vivo in an animal. Thus, in one aspect of the invention, the mixture is contained within a eukaryotic cell and the method of the invention may be performed wherein some of the components of the assay mixture may be provided exogenously to a cell by microinjection of the components therein, and some of the  
20 components may be endogenous in the cell.

The term "endogenous in the cell" as used herein, means that the component is naturally produced in the subject cell.

The term "exogenous to the cell" as used herein, means that the component is not found naturally in the subject cell, or is found therein at a low level, and is added thereto.

25 When the method of the invention is performed using a eukaryotic cell, one or more of the kinase protein, the kinase substrate and the test compound may be injected into the eukaryotic cell prior to the incubation. The cell so injected is then incubated under conditions that facilitate protein kinase activity and the level of protein kinase activity is subsequently measured following the incubation period using the assays described  
30 herein.

The eukaryotic cell that is useful in the methods of the invention may be any one of a *Xenopus laevis* oocyte, a *Xenopus laevis* embryo cell, a mammalian cell (such as a I OTI/2 cell), a *Drosophila melanogaster* S2 cell, a *Dictyostefium discoideum* cell and a yeast cell. Still more preferably, the eukaryotic cell is the murine pituitary corticotroph-  
35 derived adenoma cell line cell AtT-20.

The source of phosphate for use in the methods of the invention may be any common source of phosphate, including, but not limited to, a nucleotide triphosphates such as,

-16-

but not limited to, ATP or GTP. In a preferred embodiment, the phosphate source has bound thereon a detectable label which label is transferred with the phosphate group to the kinase substrate during the reaction. In this manner, phosphorylated kinase substrate may be distinguished from non- phosphorylated kinase substrate in that the

5 phosphorylated substrate will contain the detectable label whereas the non-phosphorylated substrate will not contain the label. In another embodiment, the phosphate source does not have bound thereon a detectable label; instead, phosphorylated kinase substrate may be distinguished from non- phosphorylated kinase substrate, for instance by recognition of one form of the substrate, but not the other, by  
10 an antibody.

The detectable label which is useful in the methods of the invention may include any known or heretofore unknown detectable label which is transferred to the kinase substrate upon transfer of a phosphate group thereto as a result of protein kinase activity. Labels which are useful include, but are not limited to, radioactive labels, such  
15 as  $\gamma^{32}\text{P}$ ,  $^{31}\text{S}$ , and non-radioactive labels, such as biotin and the like.

Preferably the kinase assay is performed in a cell and comprising;

a) contacting a cell comprising a kinase selected from the group consisting of  
SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID  
NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID  
20 NO.37, SEQ ID No.39 and SEQ ID No.41, with a source of phosphate and a suitable kinase substrate;

b) incubating said mixture in the presence or absence of a said compound and ;

c) compare the CRH response activity of said cell in the presence and absence  
of said compound.

25

Similarly, a change in CRH signaling response may be assessed using a phosphatase assay, said assay comprising;

a) contacting a mixture comprising a phosphatase selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID  
30 NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41, and a suitable phosphorylated substrate;

b) incubating said mixture in the presence or absence of a said compound and ;

measuring the level of phosphorylation of said substrate in the presence of said  
35 compound compared to the level of phosphorylation of said substrate in the absence of said test compound.

-17-

As for the kinase assay, the phosphatase in the assay of the invention, may be provided as a protein or it may be provided in the assay mixture as an mRNA encoding said phosphatase. The phosphorylated substrate is typically labeled with a detectable phosphate residue. Labels which are useful include, but are not limited to, radioactive labels, such as  $\gamma^{32}\text{P}$ ,  $^{31}\text{S}$ , and non-radioactive labels, such as biotin and the like. For use in a phosphatase activity assay, the substrate preferably consists of a peptide substrate, phosphorylated at a tyrosine or serine residue, typically labeled with  $\gamma^{32}\text{P}$ . In general, phosphorylation may be accomplished in a variety of ways. Typically, a protein tyrosine kinase is used. For example, a soluble EGF-receptor kinase in combination with  $\gamma^{32}\text{P}$ -labeled ATP may be used to phosphorylate a tyrosine residue on a peptide of the present invention. Such a phosphorylation reaction is typically allowed to proceed for about 2 hours at 30.degree. C. or overnight at room temperature. Phosphorylated peptide, hereinafter referred to as "phosphopeptide", is then purified from a phosphorylation reaction mixture. For example, peptide may be separated from a reaction mixture by addition of trichloroacetic acid and centrifugation, whereby the peptide remains in the supernate. The peptide is generally further purified by column chromatography, e.g., on C18. Purified phosphorylated peptide may be lyophilized and stored at - 20.degree. C. prior to use.

Following incubation, phosphopeptide which is not dephosphorylated ("non-dephosphorylated phosphopeptide") is separated from radioactivity released by dephosphorylation of phosphopeptide (i.e., from free radioactive phosphorus released by dephosphorylation). As used herein, the term "radioactive phosphorous" includes all forms in which a radioactive phosphorus atom may be present on a tyrosine residue and removed by dephosphorylation, e.g., as a phosphate group. Typically, separation of non-dephosphorylated phosphopeptide from free radioactive phosphorus released by dephosphorylation of phosphopeptide is effected by centrifugation, following termination of the dephosphorylation reaction by the addition of substances including nonradioactive phosphates and charcoal. Radioactivity in the supernate is determined by means well known to those of ordinary skill in the art. Based upon the amount of radioactivity added to the assay mixture initially via the phosphopeptide and the amount of radioactivity detected at the end of the assay as radioactivity released by dephosphorylation, the phosphatase enzymatic activity of the sample assayed may be calculated.

Preferably the phosphatase assay is performed in a cell and comprising:

- a) contacting a cell comprising a phosphatase selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ



-18-

ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41, with a suitable phosphorylated substrate;

b) incubating said mixture in the presence or absence of a said compound and ;

5 c) compare the CRH response activity of said cell in the presence and absence of said compound, wherein the CRH response activity of said cell is determined by the change in phosphorylation level of the substrate.

10 It is also an embodiment of the present invention to provide a method for identifying a compound capable to alter the CRH signalling response activity in a cell, said method comprising;

a) contacting a cell which expresses at least one protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID 15 No.41 with said test compound; and

b) compare the levels of a second messenger, such as cAMP, cGMP,  $Ca^{2+}$  or  $IP_3$  in said cell, in the presence and absence of said compound.

Levels of second messengers can be determined using art known techniques either in whole cells or cellular extracts comprising one of the aforementioned proteins.

20

A further method to identify a compound capable to alter CRH signalling in a cell is based on the use of a gene, such as a reporter gene, operably linked to a gene promoter or regulatory sequence element thereof characterized in that said gene promoter or regulatory sequence element comprises a transcription factor binding site, wherein said 25 transcription factor is capable of modulating CRH signalling in a cell. In a preferred embodiment the transcription factor capable of modulating CRH signalling in a cell is being selected from SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41

30 Accordingly, the present invention provides a recombinant DNA molecule comprising the gene promoter region as defined above. In the said recombinant DNA molecule, the promoter region can be operably linked to a nucleic acid molecule encoding a detectable product, such as a reporter gene. The term "operably linked", as used herein, means functionally fusing a promoter with a gene in the proper frame to express the 35 gene under control of the promoter. As used herein, the term "reporter gene" means a gene encoding a gene product that can be identified using simple, inexpensive methods

-19-

or reagents and that can be operably linked to the promoter region or an active fragment thereof. Reporter genes such as, for example, a firefly luciferase,  $\beta$ -galactosidase, alkaline phosphatase, the bacterial chloramphenicol acetyl transferase or green fluorescent protein reporter gene, can be used to determine transcriptional activity in screening assays according to the invention (see, for example, Goeddel (ed.), *Methods Enzymol.*, Vol. 185, San Diego: Academic Press, Inc. (1990); see also Sambrook, *supra*). In a preferred embodiment, the reporter gene is the firefly luciferase gene. The invention also provides a vector comprising the recombinant DNA molecule as defined above, as well as a host cell stably transformed with such a vector, or generally with the recombinant DNA molecule according to the invention. The term "vector" refers to any carrier of exogenous DNA that is useful for transferring the DNA into a host cell for replication and/or appropriate expression of the exogenous DNA by the host cell. Accordingly, in a specific embodiment said vector is an expression vector such as the commercially available pGL3luc expression vector.

15

*Identification of a CRH induced gene expression profile.*

In another aspect the present invention relates to isolated and purified nucleic acid molecules which were shown to be either up or down-regulated upon prolonged exposure to elevated levels of CRH, wherein said nucleic acid molecule is either RNA, DNA, cDNA or genomic DNA.

In particular, the present invention encompasses an isolated and purified nucleic acid molecule comprising a member selected from a group consisting of:

- a) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to a polynucleotide selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID No.5, SEQ ID No.6, SEQ ID No.7, SEQ ID No.8, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30 and SEQ ID No.31, over the entire length of said polynucleotide;
- b) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity,

even more preferably at least 97-99% identity, to a polynucleotide selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID No.5, SEQ ID No.6, SEQ ID No.7, SEQ ID No.8, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30 and SEQ ID No.31, over the entire coding region of said polynucleotide;

c) a nucleotide sequence polynucleotide selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID No.5, SEQ ID No.6, SEQ ID No.7, SEQ ID No.8, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30 and SEQ ID No.31;

Identity or similarity, as known in the art, are relationships between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. Both identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity and similarity between two polynucleotide or two polypeptide sequences, both terms are well known to skilled artisans (Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., (1988) SIAM J. Applied Math., 48, 1073. Methods commonly employed to determine identity or similarity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., (1988) SIAM J. Applied Math., 48, 1073. Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., (1984) Nucleic Acids Research 12(1), 387), BLASTP, BLASTN, and FASTA (Atschul, S. F. et al., (1990) J. Molec. Biol. 215, 403).

The nucleic acid sequence encoding a protein capable of modulating CRH activity, or fragment thereof, can be isolated from a tissue in which said gene is

expressed, such as but not limited to, brain, heart, kidney, pancreas, liver and skin. Said sequence can also be isolated from mammals other than human and mouse. Other cells and cell lines may also be suitable for use to isolate mammalian purine permease cDNA. Selection of suitable cells may be done by screening for CRH  
5 modulating activity in cell extracts or in whole cell assays, as described herein. Cells that possess CRH modulating activity in any one of these assays may be suitable for the isolation of purine permease DNA or mRNA.

Any of a variety of procedures known in the art may be used to molecularly clone DNA encoding a protein according to the invention. In one method, mRNA  
10 is isolated, and first strand cDNA synthesis is carried out. A second round of DNA synthesis can be carried out for the production of the second strand. Subsequently by the specific PCR amplification of DNA fragments through the design of degenerate oligonucleotide primers from the amino acid sequence of the purified protein that modulates CRH signalling, an isolated cDNA can be obtained. If  
15 desired the double-stranded cDNA can be cloned into any suitable vector, for example, a plasmid, thereby forming a cDNA library. Another method is to screen cDNA libraries constructed in a bacteriophage or plasmid shuttle vector with a labeled oligonucleotide probe targeted to any suitable region of SEQ ID NO: 1, SEQ ID NO: 7 or SEQ ID NO: 3. See e.g. PCR Protocols: A Guide to Method and  
20 Application, Ed. M. Innis *et al.*, Academic Press (1990).

Methods for constructing cDNA libraries in a suitable vector such as a plasmid or phage for propagation in prokaryotic or eukaryotic cells are well known to those skilled in the art. [See e.g. Maniatis *et al. Supra*]. Suitable cloning vectors are well known and are widely available.

25 It is readily apparent to those skilled in the art that other types of libraries, as well as libraries constructed from other cells or cell types, may be useful for isolating the nucleic acid sequences according to the invention. Other types of libraries include, but are not limited to, cDNA libraries derived from other cells, from organisms other than human and mouse, and genomic DNA libraries that  
30 include YAC (yeast artificial chromosome) and cosmid libraries. Construction of genomic DNA libraries can be performed by standard techniques well known in the art. Well known genomic DNA library construction techniques can be found in T. Maniatis *et al. Molecular Cloning: A Laboratory Manual*, 2d Ed. Chap. 14 (1989).

The skilled artisan will appreciate that, in many cases, an isolated cDNA  
35 sequence will be incomplete, in that the region coding for the polypeptide is short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low 'processivity' (a measure of the ability of the enzyme to remain

attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during the 1<sup>st</sup> strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (Frohman et al., 1988, PNAS USA 85, 8998-9002), or recent modifications of this technique, exemplified by the Marathon<sup>TM</sup> technology (Clontech Laboratories Inc.).

#### 10 *polypeptides*

In a further embodiment this invention relates to a polypeptide in a substantially pure form which modulate CRH signalling wherein said polypeptide is encoded by an isolated and purified nucleic acid molecule according to the invention. In a preferred embodiment the polypeptide has the amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41 and functional analogs thereof

The protein according to the invention includes all possible amino acid variants encoded by the nucleic acid according to the invention including a polypeptide encoded by said molecule and having conservative amino acid changes.

Those skilled in the art will recognize that the protein which modulate CRH signalling could be obtained by a plurality of recombinant DNA techniques including, for example, hybridization, polymerase chain reaction (PCR) amplification, or *de novo* DNA synthesis (See e.g., T. Maniatis *et al.* Molecular Cloning: A Laboratory Manual, 2d Ed. Chap. 14 (1989)).

Purified biologically active protein that modulates CRH signalings may have several different physical forms. The polypeptides according to the invention may exist as full-length nascent or unprocessed polypeptides, or as partially processed polypeptides or combinations of processed polypeptides. The full-length nascent polypeptide may be post-translationally modified, amongst other, by specific proteolytic cleavage events that result in the formation of fragments of the full-length nascent polypeptide. A fragment, or physical association of fragments may have the full biological activity associated with proteins according to the invention; however, the degree of CRH modulating activity may vary between individual fragments.

Also preferred in this aspect of the invention are fragments characterized by structural or functional attributes of the polypeptide. Preferred embodiments of the

-23-

invention in this regard include fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, high antigenic index regions of the polypeptide of the invention, and combinations of such fragments. Preferred regions are those that mediate activities of the polypeptides of the invention. Most highly preferred in this regard are fragments that have a chemical, biological or other activity of the response regulator polypeptide of the invention, including those with a similar activity or an improved activity, or with a decreased undesirable activity.

*Recombinant expression of polynucleotides encoding protein which modulate CRH activity*

5 In another embodiment polynucleotides according to the invention may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce a protein that modulates CRH signalling. Techniques for such manipulations are fully described  
10 in Maniatis, T, et al., *supra*, and are well known in the art.

Therefore, in a further aspect this invention provides an expression vector for expression of a protein that modulates CRH signalling in a recombinant host, wherein said vector contains a nucleic acid sequence encoding a protein that modulates CRH signalling and functional analogs thereof. In a more preferred aspect of this invention  
15 this expression vector contains a nucleic acid molecule encoding a protein that modulates CRH signalling, having a nucleotide sequence selected from a group consisting of: SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40 and functional analogs  
20 thereof or contains genomic DNA encoding a protein that modulates CRH signalling.

Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express eukaryotic genes in a variety of hosts such as bacteria including *E. coli*, cyanobacteria, plant cells, insect  
25 cells, amphibian cells, fungal cells including yeast cells, and animal cells.

Specifically designed vectors allow the shuttling of DNA between hosts such as bacteria-yeast or bacteria-animal cells or bacteria-fungal cells or bacteria-invertebrate cells. An appropriately constructed expression vector may contain: an origin of replication for autonomous replication in host cells, selectable markers, a  
30 limited number of useful restriction enzyme sites, a potential for high copy number, and active promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and initiate RNA synthesis. A strong promoter is one that causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors,  
35 specifically designed plasmids or viruses.

The isolated and purified nucleic acid molecules, according to the invention, encoding a protein which modulates CRH signalling may be cloned into an expression vector for expression in a recombinant host cell. Recombinant host cells may be prokaryotic or eukaryotic, including but not limited to bacteria such as E. coli, fungal cells such as yeast, amphibian cells such as Xenopus oocytes, mammalian cells including but not limited to cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including but not limited to Drosophila- and silkworm-derived cell lines. Cell lines derived from mammalian species which may be suitable and which are commercially available, include but are not limited to, CV-1 (ATCC CCL 70), COS-1 (ATCC CRL 1650), COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCC CCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26), MRC-5 (ATCC CCL 171), L-cells, neuroblastoma, glial cells and HEK-293 (ATCC CRL1573).

Therefore, in a further embodiment this invention relates to a recombinant host cell containing a recombinantly cloned nucleic acid molecule encoding a protein that modulates CRH signalling or functional analog thereof. In a further aspect the recombinant host cell according to the invention contains a nucleic acid molecule which is either genomic DNA or has a nucleotide sequence selected from a group consisting of: SEQ ID NO.9, SEQ ID NO.11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID NO.38 and SEQ ID NO.40; and functional analogs thereof.

The expression vector may be introduced into host cells via any one of a number of techniques including but not limited to transformation, transfection, protoplast fusion, lipofection, and electroporation. The expression vector-containing cells are clonally propagated and analyzed to determine whether they produce a protein that modulates CRH signalling. Identification of permeases expressing host cell clones may be done by several means, including but not limited to immunological reactivity with antibodies directed against the polypeptides according to the invention, and the presence of host cell-associated mammalian purine permease activity.

Thus, the present invention also relates to a process for expression of protein that modulates CRH signalling in a recombinant host cell, comprising culturing the host cells according to the invention under conditions which allow expression of the protein that modulates CRH signalling from the expression vector as outlined herein. The proteins of this invention may be synthesized either by direct



-26-

expression or as a fusion protein comprising the protein of interest as a translational fusion with another protein or peptide that may be removable by self, enzymatic or chemical cleavage. Therefore, in a particular embodiment this invention provides the proteins according to the invention wherein said polypeptides are part of a fusion protein.

5 It is often observed in the production of certain peptides in recombinant systems that expression as a fusion protein prolongs the life span, increases the yield of the desired peptide, or provides a convenient means of purifying the protein. This is particularly relevant when expressing mammalian proteins in  
10 prokaryotic hosts. A variety of peptidases (e.g. enterokinase and thrombin), which cleave a polypeptide at specific sites or digest, the peptides from the amino or carboxy termini (e.g. diaminopeptidase) of the peptide chain are known. Furthermore, particular chemicals (e.g. cyanogen bromide) will cleave a polypeptide chain at specific sites. The skilled artisan will appreciate the  
15 modifications necessary to the amino acid sequence (and synthetic or semi-synthetic coding sequence if recombinant means are employed) to incorporate site-specific internal cleavage sites. See e.g., P.Carter, "Site Specific Proteolysis of Fusion Proteins", Chapter 13, in Protein Purification: From Molecular Mechanisms to Large Scale Processes, American Chemical Society, Washington, D.C. (1990).

20 Furthermore, one could use, e.g., a mammalian cell that already comprises in its genome a nucleic acid molecule encoding a protein that modulates CRH signalling as described above, but does not express the same or not in an appropriate manner due to, e.g., a weak promoter, and introduce into the mammalian cell a regulatory sequence such as a strong promoter in close proximity to the endogenous  
25 nucleic acid molecule encoding said purine permease polypeptide so as to induce expression of the same.

As such a recombinant host cell containing a polynucleotide encoding a protein which modulates CRH signalling under the control of a heterologous transcription and/or regulatory sequence or protein, would be another embodiment of  
30 this invention.

In this context the term "regulatory sequence" denotes a nucleic acid molecule that can be used to increase the expression of the purine permease polypeptide, due to its integration into the genome of a cell in close proximity to the CRH modulating protein-encoding gene. Such regulatory sequences comprise  
35 promoters, enhancers, inactivated silencer intron sequences, 3'UTR and/or 5'UTR coding regions, protein and/or RNA stabilizing elements, nucleic acid molecules encoding a regulatory protein, e.g., a transcription factor, capable of inducing or

triggering the expression of the CRH modulating protein-encoding gene or other gene expression control elements which are known to activate gene expression and/or increase the amount of the gene product. The introduction of said regulatory sequence leads to increase and/or induction of expression of polypeptides, which  
5 modulate CRH signalling, resulting in the end in an increased amount of said polypeptides in the cell. Thus, the present invention is aiming at providing *de novo* and/or increased expression of polypeptides that modulate CRH signalling.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods.  
10 Such methods are described in many standard laboratory manuals, such as Davis, Basic Methods In Molecular Biology (1986). It is specifically contemplated that polypeptides, which modulate CRH signalling may in fact be expressed by a host cell lacking a recombinant vector.

15 In addition, expression of polynucleotides according to the invention may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA or mRNA isolated from cells capable of modulating CRH signaling can be efficiently translated in various cell-free systems, including but not limited to wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based  
20 systems, including but not limited to microinjection into frog oocytes, with microinjection into frog oocytes being generally preferred.

#### *Transgenic non-human animals*

The present invention also relates to a method for the production of a  
25 transgenic non-human animal, preferably transgenic mouse, comprising introduction of a polynucleotide or vector of the invention into a germ cell, an embryonic cell, stem cell or an egg or a cell derived therefrom. The non-human animal can be used in accordance with a screening method of the invention described herein and may be a non-transgenic healthy animal, or may have a phosphate uptake or reabsorption  
30 disorder, preferably a disorder caused by at least one mutation in the protein that modulates CRH signalling. Such transgenic animals are well suited for, e.g., pharmacological studies of drugs in connection with mutant forms of the above described polypeptides. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., Gene Targeting, A Practical  
35 Approach (1993), Oxford University Press. The DNA of the embryonal membranes

of embryos can be analyzed using, e.g., Southern blots with an appropriate probe; see supra.

Preferably, the transgenic non-human animal of the invention further comprises at least one inactivated wild type allele of the corresponding mammalian CRH modulating protein-encoding gene; see supra. This embodiment allows for example, the study of the interaction of various mutant forms of polypeptides according to the invention on the onset of the clinical symptoms of disease related to disorders in CRH metabolism. All the applications that have been herein before discussed with regard to a transgenic animal also apply to animals carrying two, three or more transgenes; e.g. encoding neutral endopeptidase (NEP). It might be also desirable to inactivate protein that modulates CRH signalling expression or function at a certain stage of development and/or life of the transgenic animal. This can be achieved by using, for example, tissue specific, developmental and/or cell regulated and/or inducible promoters which drive the expression of, e.g., an antisense or ribozyme directed against the RNA transcript encoding the protein capable of modulating CRH signalling; see also supra. A suitable inducible system is for example tetracycline-regulated gene expression as described, e.g., by Gossen and Bujard (Proc. Natl. Acad. Sci. 89 USA (1992), 5547-5551) and Gossen et al. (Trends Biotech. 12 (1994), 58-62). Similar, the expression of the mutant protein that modulates CRH signalling may be controlled by such regulatory elements.

Furthermore, the invention also relates to a transgenic mammalian cell which contains (preferably stably integrated into its genome) a nucleic acid molecule according to the invention or part thereof, wherein the transcription and/or expression of the nucleic acid molecule or part thereof leads to reduction of the synthesis of a protein that modulates CRH signalling.

In a preferred embodiment, the reduction is achieved by an anti-sense, sense, ribozyme, co-suppression and/or dominant mutant effect. "Antisense" and "antisense nucleotides" means DNA or RNA constructs which block the expression of the naturally occurring gene product.

The provision of the polynucleotide according to the invention opens up the possibility to produce transgenic non-human animals with a reduced level of the protein as described above and, thus, with a defect in phosphate metabolism. Techniques how to achieve this are well known to the person skilled in the art. These include, for example, the expression of antisense-RNA, ribozymes, of molecules which combine antisense and ribozyme functions and/or of molecules

which provide for a co-suppression effect; see also supra. When using the antisense approach for reduction of the amount of protein that modulates CRH signalling in cells, the nucleic acid molecule encoding the antisense-RNA is preferably of homologous origin with respect to the animal species used for transformation. However, it is also possible to use nucleic acid molecules which display a high degree of homology to endogenously occurring nucleic acid molecules encoding a protein that modulates CRH signalling. In this case the homology is preferably higher than 80%, particularly higher than 90% and still more preferably higher than 95%. The reduction of the synthesis of a protein according to the invention in the transgenic mammalian cells can result in an alteration in, e.g., adenine reabsorption. In transgenic animals comprising such cells this can lead to various physiological, developmental and/or morphological changes.

Thus, the present invention also relates to transgenic non-human animals comprising the above-described transgenic cells. These may show, for example, a deficiency in CRH metabolism compared to wild type animals due to the stable or transient presence of a foreign DNA resulting in at least one of the following features:

- (a) disruption of (an) endogenous gene(s) encoding a protein capable of modulating CRH signalling;
- (b) expression of at least one antisense RNA and/or ribozyme against a transcript comprising a polynucleotide of the invention;
- (c) expression of a sense and/or non-translatable mRNA of the polynucleotide of the invention;
- (d) expression of an antibody of the invention;
- (e) incorporation of a functional or non-functional copy of the regulatory sequence of the invention; or
- (f) incorporation of a recombinant DNA molecule or vector of the invention.

With the polypeptides, their encoding polynucleotides and vectors of the invention, it is now possible to study *in vivo* and *in vitro* the efficiency of drugs in relation to particular mutations in protein that modulates CRH signalling of a patient and the affected phenotype. Furthermore, mutant forms of polypeptides of the invention can be used to determine the pharmacological profile of drugs and for the identification and preparation of further drugs which may be effective for the treatment of disorders related to the CRH metabolism, in particular for the amelioration of CRH induced stress or depression.

It will thus be appreciated that the present invention also relates to a method for preventing, treating or ameliorating a medical condition related to a disorder of CRH metabolism including CRH receptor related disorders which comprises  
5 administering to a mammalian subject a therapeutically effective amount of the polypeptides, the polynucleotides or the vectors encoding a protein capable of modulating CRH signalling of the present invention.

#### *Diagnostic Assays*

10 This invention further relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ  
15 ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40 which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or  
20 altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

It will thus be appreciated that this invention provides a method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to a disorder of CRH activity comprising:

25 (a) determining the presence or absence of a mutation in the polynucleotide according to the invention; and

(b) diagnosing a pathological condition or susceptibility to a pathological condition based on the presence or absence of said mutation.

Nucleic acids for diagnosis may be obtained from a subject's cells, such as from  
30 blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be  
35 identified by hybridizing amplified DNA to labeled nucleotide sequences according to the invention. Perfectly matched sequences can be distinguished from mismatched

duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in capillary electrophoresis columns or gels, with or without denaturing agents, or by direct DNA sequencing (e.g., Myers *et al.*, *Science* (1985)230:1242). Sequence changes at specific locations may also be revealed by specific restriction endonucleases, nuclease protection assays, such as RNase and S1 protection or a chemical cleavage method (see Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85: 4397-4401). In another embodiment, an array of oligonucleotides probes comprising a nucleotide sequence encoding a protein capable of modulating CRH activity or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

The diagnostic assays offer a process for diagnosing or determining a susceptibility to the diseases through detection of mutations in the CRH modulating protein-encoding gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of polypeptide or mRNA, as well as by determining from said samples the presence of protein derivatives compared to the normal structure (see above). Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example; nucleic acid amplification, for instance via PCR, RT-PCR; RNase protection; Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Assay techniques that can be used to determine the presence of protein derivatives or variants comprise amongst others mass spectrometry.

Thus in another aspect, the present invention provides a method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to a disorder of prolonged CRH exposure comprising:

- (a) determining the presence or amount of expression of the polypeptide or a derivative thereof according to the invention in a biological sample; and
- (b) diagnosing a pathological condition or susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide or of a

derivative thereof.

In particular the present invention provides a method of diagnosing a CRH induced gene expression profile in an individual, said method comprising;

a) obtaining a biological sample of said individual; and

5 b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said biological sample;

whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID 10 NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

Preferably the amount of said proteins is determined either at the protein level, preferably using antibodies that bind thereto, or at the gene transcription level, preferably using probes that bind to a polynucleotide encoding an amino acid 15 sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

Alternatively the CRH induced gene expression profile is determined by assessing 20 the level of gene transcription of a gene comprising a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID 25 No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40. Methods to determine the level of gene transcription have been described hereinbefore and comprise in a preferred embodiment the use of a probe which binds, preferably selectively binds to a polynucleotide selected from the group consisting of SEQ ID No.1, SEQ ID No.2, 30 SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40 or the 35 complement thereof. In another embodiment, an array of oligonucleotides probes comprising a nucleotide sequence according to the invention or fragments thereof

can be constructed to conduct efficient screening of the level of gene transcription in the sample of an individual.

5 In a further aspect, the present invention relates to a diagnostic kit which comprises:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID NO.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID NO.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40 or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

15 (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41 or a fragment thereof; or

20 (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41 and optionally suitable means for detection.

25 It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly CRH metabolism related disorders such as CRH induced stress or depression.

30 The nucleotide sequences of the present invention are also valuable for chromosome localisation. These sequences are specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene-associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library).  
35 The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of



physically adjacent genes). The gene of the present invention maps to human chromosome 15.

5 The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

10 The nucleotide sequences of the present invention are also valuable for tissue localisation. Such techniques allow the determination of expression patterns of the polypeptides according to the invention in tissues by detection of the mRNAs that encode them. These techniques include *in situ* hybridization techniques and nucleotide amplification techniques, for example PCR. Such techniques are well known in the art. Results from these studies provide an indication of the normal functions of the polypeptides in the organism.

15 The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them, can also be used as immunogens to produce antibodies immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other-related polypeptides in the prior art.

20 Thus in another embodiment, this invention provides a monospecific antibody immunologically reactive with a mammalian purine permease. In a preferred embodiment said antibody is immunologically reactive with a polypeptide having an amino acid sequence selected from a group consisting of: SEQ ID NO.10, SEQ ID  
25 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41; and functional analogs thereof or said antibody blocks activity of a protein that modulates CRH signalling.

30 Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells expressing these to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975)256:495-497),  
35 the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983)4:72) and the EBV-hybridoma technique (Cole *et al.*, *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, pp.77-96, Alan R.

Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those described in U.S. Patent No.4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other  
5 organisms, including other mammals, may be used to express humanized antibodies.

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

Antibodies against polypeptides of the present invention may also be employed  
10 to treat the CRH metabolism related disorders.

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgD, IgE). Preferred as  
15 an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG I, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with for instance blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion  
20 proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. W094/29458 and W094/22914.

## 25 *Therapeutic Utility*

A further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a polypeptide of the present invention wherein the composition comprises a polypeptide or polynucleotide of the present  
30 invention. The vaccine formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers,  
35 bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include

suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

In still another approach, expression of the gene encoding proteins which modulate CRH signalling can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or externally administered (see, for example, O'Connor, *J.Neurochem* (1991) 56:560 ;Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices ("triplexes") with the gene can be supplied (see, for example, Lee *et al.*, *Nucleic Acids Res* (1979)6:3073; Cooney *et al.*, *Science* (1988)241:456; Dervan *et al.*, *Science* (1991)251:1360). These oligomers can be administered per se or the relevant oligomers can be expressed *in vivo*. Synthetic antisense or triplex oligonucleotides may comprise modified bases or modified backbones. Examples of the latter include methylphosphonate, phosphorothioate or peptide nucleic acid backbones. Such backbones are incorporated in the antisense or triplex oligonucleotide in order to provide protection from degradation by nucleases and are well known in the art. Antisense and triplex molecules synthesised with these and/or other modified backbones also form part of the present invention.

In another process for inhibiting expression of a target gene in a cell, RNA with partial or fully double-stranded character is introduced into the cell or into the extracellular environment. Inhibition is specific in that a nucleotide sequence from a portion of the target gene is chosen to produce inhibitory RNA. The RNA may comprise one or more strands of polymerized ribonucleotide; it may include modifications to either the phosphate-sugar backbone or the nucleoside. The double-stranded structure may be formed by a single self-complementary RNA strand or two complementary strands. Inhibition is sequence-specific in that the nucleotide sequences corresponding to the duplex region of the RNA are targeted for genetic inhibition. RNA containing a nucleotide sequence identical to a portion of the target sequence is preferred. Examples of RNA inhibition technology can be found in International Patent Application WO 99/32619.

In addition, expression of the proteins which modulate CRH signalling may be

prevented by using ribozymes specific to the mRNA sequence encoding said protein. Ribozymes are catalytically active RNAs that can be natural or synthetic (see for example Usman, N, et al., Curr. Opin. Struct. Biol (1996)6(4), 527-33.) Synthetic ribozymes can be designed to specifically cleave the aforementioned mRNAs at selected positions thereby preventing translation of said mRNAs into functional polypeptide. Ribozymes may be synthesised with a natural ribose phosphate backbone and natural bases, as normally found in RNA molecules. Alternatively the ribozymes may be synthesised with non-natural backbones to provide protection from ribonuclease degradation, for example, 2'-O-methyl RNA, and may contain modified bases.

For treating abnormal conditions related to an under-expression of proteins which modulate CRH signalling, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates a polypeptide of the present invention, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of mammalian purine permease by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication-defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of a polypeptide of the present invention in combination with a suitable pharmaceutical carrier.

In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide, such as the soluble form of a polypeptide of the present invention, agonist/antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers

filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

5       The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using  
10       penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of patches, salves, pastes, gels, and the like.

15       The dosage range required depends on the choice of peptide or other compounds of the present invention, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages; however, are in the range of 0.1-100 $\mu$ g/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the  
20       variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

25       Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into  
30       the subject.

      This invention will be better understood by reference to the Experimental Details that follow, but those skilled in the art will readily appreciate that these are only illustrative of the invention as described more fully in the claims that follow thereafter.  
35       Additionally, throughout this application, various publications are cited. The disclosure of these publications is hereby incorporated by reference into this application to describe more fully the state of the art to which this invention pertains.

### EXPERIMENTAL PROCEDURES

5 *Animal housing and treatment* – Male transgenic animals and their wild-type littermates were maintained in an specific pathogen free facility that meets all national and European requirements for animal care. Mice were housed individually 4 weeks before the start of the experiment in a climate-controlled animal colony with a 12h dark-light cycle (light on 7:00 EST) with free access to food and water. Prior to treatment 3 month old mice were randomized over three groups each comprising 4 wild-type and 4  
10 transgenic animals. One group received no treatment, a second group received vehicle injection 5 days, 2 times per day (10% cyclodextran (CD), pH4) and a third group received 10 mg/kg of a CRH-R1 specific antagonist R121919 in 10% CD following the same regime. 16 hrs after last treatment animals were sacrificed for dissection of brain areas and subsequent isolation of RNA.

15 *Sample preparation* – Macroscopically the following brain areas were dissected from transgenic and wild-type animals; pituitary, cerebellum, temporal area, hippocampus, frontal cortex, and nucleus accumbens. Brain tissue was homogenized in Trizol (Invitrogen Life Technologies) using an Ultra-turrax T25 grinder (IKA-labortechnik). Total RNA was extracted using Trizol according to the instructions of the  
20 manufacturer. Total RNA was further purified using Rneasy kit (Qiagen) with DNaseI treatment on column.

*Microarray hybridization* – cRNA was prepared as follows. Reverse transcription was performed on 10µg of total RNA for 1h at 42°C using a T7-oligo(dT)<sub>24</sub>-primer and SuperscriptII RT (Invitrogen Life Technologies). Second strand cDNA synthesis was  
25 done for 2 h at 16°C using *Escherichia coli* DNA Polymerase I, DNA ligase and RNaseH (Invitrogen Life Technologies). After phenol-chloroform extraction using phase-lock gel (Eppendorf) in vitro transcription was performed for 6 h at 37°C using the Bioarray high-yield RNA transcript labeling kit with Biotin labeled ribonucleotides (Enzo Diagnostics). cRNA samples were purified on Qiagen Rneasy columns followed  
30 by fragmentation for 35 min at 95°C. cRNA yields were between 50 and 100µg. Samples were processed on GeneChips (Affymetrix, Santa Clara, CA). In order to check the quality of each sample, 5µg of labeled cRNA was run on Test2-arrays. Actual experiments were performed on Murine Genome U74Av2 arrays, containing probe sets interrogating approximately 12,000 full-length mouse genes and EST clusters from the  
35 UniGene database (Build 74). Hybridization was performed using 15µg cRNA for 16 h at 45°C under continuous rotation. Arrays were stained in Affymetrix Fluidics stations

- using Streptavidin/Phycoerythrin (SAPE) followed by staining with anti-streptavidin antibody and a second SAPE staining. Subsequently arrays were scanned with a HP-Laserscanner and data were analyzed with the Microarray Suite Software (Affymetrix). No scaling or normalization was performed at this stage. Quality of the experiment was assessed based on the percentages present calls (see table).

Area	Present calls	3'/5' $\beta$ -Actin	3'/5' GAPDH
temporal area	46.15 $\pm$ 1.98%	1.96 $\pm$ 0.53	1.46 $\pm$ 0.38
hippocampus	46.27 $\pm$ 1.89%	1.58 $\pm$ 0.39	1.22 $\pm$ 0.29
cerebellum	47.14 $\pm$ 2.13	1.67 $\pm$ 0.37	0.97 $\pm$ 0.08
frontal cortex	44.68 $\pm$ 2.23	1.85 $\pm$ 0.39	1.29 $\pm$ 0.26
nucleus accumbens	44.27 $\pm$ 2.11	2.69 $\pm$ 0.85	2.52 $\pm$ 1.20
pituitary	45.33 $\pm$ 1.67	1.53 $\pm$ 0.28	1.25 $\pm$ 0.21

- Samples with low yields of RNA or low percentages present calls or deviating 3'/5' ratios were omitted from subsequent analysis. The number of animals that were used for analysis are shown in table.

	WT untreated	WT vehicle	WT R121919	TG untreated	TG vehicle	TG R121919
temp. area	4	4	4	3	4	3
hippocampus	4	4	4	3	3	4
cerebellum	3	4	4	3	4	3
frontal cortex	3	4	4	3	4	4
nucleus accumbens	4	3	4	4	3	4
pituitary	4	4	4	4	3	4

## 15 Data analysis and selection of genes

Raw intensities on each array were normalized using the following algorithm:

$$\log I_{aligned} = \log I_{raw} + \left( \log \frac{\sum_{all\ samples} I_{raw}}{\# genes \times \# samples} - \log \frac{\sum_{sample} I_{raw}}{\# genes} \right)$$

Basically this alignment sets the average intensity of one array to the average measured this normalization compensates for array to array variations in hybridization, washing and staining, ultimately allowing a reasonable comparison between arrays. After  
5 normalization, data were analyzed using weighted spectral mapping analysis (SMA) (7). Weighted spectral mapping is an unsupervised multivariate analysis method which includes double-centering of the data combined with a specialized visualization representing the two highest principle components explaining most of the variance within the dataset. Even though double-centering removes the "size" component of the  
10 array data, this information is reintroduced in the visualization via the area of the symbols representing the size of the respective samples (depicted by squares) and genes (depicted by circles). What remains are contrasts between the different genes and contrasts between the different samples of the data table. These contrasts can be expressed as ratios due to logarithmic transformation performed by the algorithm. The  
15 contrasts can be understood as specificities of the different genes for the different samples. Conversely, they refer also to the specificities or preferences of the different samples for some of the genes. Therefore, one could state that SMA provides a visualization of the interactions between genes and samples. This method allows the reduction of a large microarray dataset and provides means to visually inspect and  
20 thereby identify clusters of genes and/or subjects in the data. In the spectral map samples which cluster together can thus be interpreted as being similar. Correspondingly samples located at opposite sites of the axes or the center can be interpreted as being dissimilar.

In addition to this overall analysis the identification of individual genes with a different  
25 expression level between groups were identified using significance analysis of microarray data (SAM) (8). SAM identifies genes with statistically significant changes in expression by assimilating a set of gene-specific t-tests. In contrast to the classical t-test, this gene-specific t-tests not only takes into account the variance observed for a particular gene but also includes a so called fudge factor covering the overall variance  
30 across all genes measured. Genes with scores greater than a threshold are deemed potentially significant. In order to estimate the number of false positives among these potentially significantly changed genes, SAM uses permutations to estimate the false discovery rate. To this end t-values are calculated iteratively on permutations of the measurements. The threshold can be adjusted to identify smaller or larger sets of genes,  
35 and corresponding FDRs are calculated. In addition to the threshold delta that influences the number of significantly changed genes and a corresponding FDR, a fold change threshold can be imposed to discards genes that show only a limited difference



in expression between samples. These changes are probably from a biological point of view of little significance.

K-means clustering of expression data obtained on a given brain area was carried out using the OmniViz program. All records which were absent in all samples were

- 5 removed and all signals less than 20 were set to 20. This allowed the clustering of genes with similar behavior.

**Quantitative RT-PCR** - Microarray data were confirmed using real time PCR analysis. First strand cDNA synthesis was performed on 0.5µg total RNA using random hexamer primers and SuperscriptII RT (Invitrogen Life Technologies). Quantitative PCR was  
10 performed on a ABIPrism 7700 cyclor (Applied Biosystems) using a Taqman PCR kit. Serial dilutions of cDNA were used to generate standard curves of threshold cycles versus the logarithms of concentration for  $\beta$ -actin, CRH receptor 1 (Crh-R1), CRH receptor 2 (Crh-R2), neurotensin receptor 1 (Ntsr1), neurotensin receptor 2 (Ntsr2), neurotensin receptor 3 (Ntsr3), 11 $\beta$ -hydroxysteroid dehydrogenase 1 (Hsd11B1) and  
15 serum/glucocorticoid regulated kinase (Sgk). A linear regression line calculated from the standard curves allowed the determination of transcript levels in RNA samples from the different treatment groups at each brain area.

$\beta$ -actin forward	5'-CATCTTGGCCTCACTGTCCAC-3'
$\beta$ -actin probe	5'-TGCTTGCTGATCCACATCTGCTGGA-3'
$\beta$ -actin reverse	5'-GGGCCGGACTCATCGTACT-3'
Crh-R2 forward	5'-GGGAGAACAGAAGCGCCTG-3'
Crh-R2 probe	5'-AGAAGGOTGAGGATCCCCAAATCAGAGT-3'
Crh-R2 reverse	5'-CCCTTGTTTCAATCACTCCCA-3'
Crh-R1 forward	5'-TTTCTGAACAGTGAGGTCCGC-3'
Crh-R1 probe	5'-CCGGAAGAGGTGGCGGCGA-3'
Crh-R1 reverse	5'-GGGCTCTGATGGAGTGCTTG-3'
Ntsr1 probe	5'-CCAACGTTCTCCAGGAAGCCAAACAG-3'
Ntsr1 reverse	5'-ACGCATGGTTGCTGGACAT-3'
Ntsr2 forward	5'-TGGTGACCAACACGCTCTTCT-3'
Ntsr2 probe	5'-TCAGCTCGGCAGTGACCCCA-3'
Ntsr2 reverse	5'-AGGAAGACACGGCGTTGTAGA-3'
Ntsr3 forward	5'-GGAAGCCGGAGAACAGCAA-3'
Ntsr3 probe	5'-TGCGACGCTACCGCAAAGAACA-3'
Ntsr3 reverse	5'-GGATATGAAGGCTGCACTCGTT-3'
Sgk forward	5'-TGGACCAATGCCCCAGTT-3'

Sgk probe	5'-TCAGTCAAAGCCGTTGGTGTTCATTG-3'
Sgk reverse	5'-GCCCCGTTTTATAGGTGACATTTTAA-3'

### RESULTS & DISCUSSION

In order to study the effects of prolonged CRH administration on the central nervous system we compared gene expression profiles in brain regions derived from CRH overexpressing mice to those of their wild-type littermates. In addition we studied the effects of CRH-R1 antagonist administration on these expression profiles. The brain regions investigated in this study include pituitary, nucleus accumbens, frontal cortex, temporal area, hippocampus and cerebellum. Expression of CRH receptor 1 (CRH-R1) and 2 (CRH-R2) was measured in brain areas using quantitative real time RT-PCR (RTq). Although CRH-R1 was readily detectable, no reliable measurements could be obtained for CRH-R2 in these brain areas. The observed expression levels of CRH receptors in these brain areas correspond with previous reports of high levels of CRH-R1 in cerebellum and frontal cortex and lower levels of expression of CRH-R2 which is mainly a peripheral receptor (see figure 1)(9).

RNA isolated from the different brain areas derived from different animals were processed separately on arrays containing 12000 murine EST. After normalization data were analyzed per brain area. First pass examination of data was performed using spectral map analysis (SMA), allowing an overall graphical interpretation of gene expression responses in relation to the treatment group and genotype. SMA indicated no gross changes in expression between treatment groups for all areas except pituitary. In all brain areas most samples derived from transgenic animals (Tg) are clustered at a different location than those derived from wild-type animals (WT). This is most obvious at the level of the pituitary where a clear difference in expression pattern is observed between Tg and WT animals with samples clustered at opposite sites in the spectral map. Taken together these data indicate that life long exposure to CRH induces changes in gene expression profiles in all studied brain areas, with the most prominent effects observed in pituitary. These changes in expression profile are not affected (or only to a limited degree) by a 5 day treatment with CRH-R1 antagonist as exemplified by the pituitary spectral map.

In order to identify genes that were significantly changed, the significance analysis of microarray data (SAM) algorithm was applied. In agreement with the SMA only a limited number of genes was found significantly changed in most brain areas (see supplementary information). In pituitary however more than 300 genes were differently

expressed between WT and TG animals. Gene expression profiles in the studied brain areas of CRH overexpressing mice elucidated previously unrecognized homeostatic mechanisms present in these animals. In general the observed expression profiles point towards a few pathways that are affected in transgenic animals (TG).

- 5 TG animals have plasma glucocorticoid levels that are 6-8 times higher than in WT animals (see figure 3). As a consequence gene expression profiles show an adaptation to these high levels of corticosterone. This is achieved both by downregulation of enhancers of glucocorticoid receptor signaling such as 11 $\beta$ -hydroxysteroid dehydrogenase type 1 (11 $\beta$ -HSD1) in hippocampus and upregulation of putative  
10 repressors of this signaling pathway such as FK506 binding protein 5 (Fkbp5) (see figure 4).

Tissue glucocorticoid concentrations are determined not only by plasma hormone levels, but also by two intracellular 11 $\beta$ -hydroxysteroid dehydrogenases (type 1 and 2), which locally interconvert active glucocorticoids and inert 11-keto forms. 11 $\beta$ -HSD1,  
15 the predominant isoform in the brain, appears to function as predominant 11 $\beta$ -reductase, regenerating active glucocorticoids from 11-keto forms. By studying 11 $\beta$ -HSD1 deficient mice it was demonstrated that the intracellular regeneration of glucocorticoids by 11 $\beta$ -HSD1 plays an important role in HPA-axis control. In response to restraint stress these mice show exaggerated ACTH and corticosterone levels. In  
20 addition 11 $\beta$ -HSD1 deficient mice were less sensitive to exogenous cortisol suppression of HPA activation, suggesting a diminished glucocorticoid feedback in these animals. In view of the significant downregulation of 11 $\beta$ -HSD1 in hippocampus of CRH overexpressors it can be envisaged that these mice have an altered glucocorticoid feedback.

- 25 Fbp5 was found upregulated in all brain areas tested, although only significant in a few, all other showed a similar tendency (figure 4). The observation that the exchange of the immunophilin Fkbp5 for Fkbp4 is an important first step of for activation of the glucocorticoid receptor (GR) is of interest. In squirrel monkey and New World primate genera in general, Fkbp5 has been identified a potent inhibitor of glucocorticoid  
30 receptor binding. Our observation of induction of Fkbp5 could be explained as an attenuation of the GR in response persistent high levels of circulating glucocorticoid.

Another indication of altered glucocorticoid signaling is the upregulation of serum/glucocorticoid kinase (Sgk) in cerebellum, nucleus accumbens and temporal area (figure 4). Sgk is a serine/threonine protein kinase of which transcription is induced by  
35 glucocorticoid and serum and is regulated by anisotonic and isotonic cell volume changes. In rat brain it has been shown that Sgk mRNA levels were increased by dehydration in the temporal lobe, including hippocampus. In the same study it was

demonstrated that Sgk markedly increased the activity of the neuronal  $K^+$  channel Kv1.3 suggesting that Sgk might participate in the regulation of neuronal excitability. Of interest is a recent report on facilitation by Sgk of memory consolidation of spatial learning in rats. Sgk was shown to be differently expressed in fast learners versus slow learners, moreover it was demonstrated that transfection of Sgk into the CA1 region of hippocampus leads to increased water maze performance (10). In that respect it is striking that while this study demonstrates that CRH overexpressors show high levels of Sgk mRNA, these animals have been reported to exhibit a profound learning deficit (11).

Neurotensin receptor 2 (Ntsr2) was found downregulated in several brain tissues including hippocampus and nucleus accumbens except pituitary (see figure 5). Neurotensin (NT) is a tridecapeptide which is found in high concentration in numerous areas of the CNS where it plays a role as neurotransmitter and neuromodulator. When NT is administered by intracerebroventricular injection a variety of CNS effects are observed, including hypolocomotion, hypothermia, analgesia, and reduced food consumption. Three receptors for NT have been identified (Ntsr1, 2 and 3). Studies of Ntsr1 deficient mice demonstrated this receptor to be responsible for the effects of NT on body temperature, feeding behavior and hypolocomotion. Ntsr2 on the other hand seems to account for the analgesic effects of NT as demonstrated by applying an antisense strategy in mice brain. The observation of Ntsr2 downregulation prompted us to investigate expression levels of other members of the NT receptor family by RTq. Ntsr3 did not show any change in expression in all areas tested. In contrast Ntsr1 was significantly downregulated in CRH overexpressors with the effect being the most pronounced in hippocampus (figure 5). Level of downregulation was higher for Ntsr1 than for Ntsr2. Striking was the upregulation of both Ntsr1 and Ntsr2 in vehicle treated animals (both tg and Wt) compared to untreated animals. This increase in expression level was abolished in R121919 treated animals, suggesting that repeated stress induced expression of Ntsr1 (and to a lesser extent Ntsr2) and that this induction could be reverted by a CRH-R1 antagonist (even 16 hrs after last treatment). Our data demonstrate for the first time a link between CRH and NT systems in CRH overexpressing animals and suggest a role for these NT receptors in repeated stress. Other pathways affected include intracellular calcium signaling/sensing (hippocalcin like 1, calcyclin, ...), myelination (myelin, myelin-associated glycoprotein, ...), cell proliferation and neurogenesis (Edg2, Id2, GAB1, ...) and extracellular matrix formation (decorin, brevican, ...).

Hippocalcin like 1 (Hpcal1) belongs to the neuronal calcium sensor family family of  $Ca^{2+}$ -binding proteins which play a role in diverse processes, including modulation of

neurotransmitter release, control of cyclic nucleotide metabolism, biosynthesis of phosphoinositides and indirect regulation of ion channels. It has been demonstrated that hippocalcin might potentiate the protection from apoptosis exerted by neuronal inhibitor-of-apoptosis (IAPs) proteins.

- 5 The changes of expression observed in genes encoding proteins involved in myelination, cell proliferation and extracellular matrix formation suggest a change in the dynamics of neurogenesis in CRH overexpressors. In support of this are the changes in expression observed mainly in nucleus accumbens involving the endothelial differentiation lysophosphatidic acid G-protein-coupled receptor 2 (Edg2), growth  
10 factor receptor bound protein 2-associated protein 1 (Gab1), Inhibitor of DNA binding 2 (Id2) and fibroblast growth factor receptor 2 (Fgfr2).

- As the pituitary is the primary target organ for CRH it is expected that prolonged elevated levels of CRH as present in CRH expressors induce major changes in expression profiles in the pituitary of transgenic animals. The activated state of the  
15 pituitary is exemplified by high expression levels of several kallikreins, which play a role as proprotein converting enzymes in the processing of diverse hormones secreted from the pituitary. Striking are the elevated levels of preproenkephalin A (10 times higher in Tg versus Wt) and to a lesser extent of prodynorphin (2 times higher in Tg versus Wt). This finding of elevated levels of endogenous opioids in CRH  
20 overexpressors is in line with the notion that these opioids represent a major modulatory system in the adaptation of an organism to chronic stress, balancing the response that the stressor places on the brain with the potentially detrimental effects that a sustained stress response may produce. Other interesting findings in the pituitary include elevation of Bdnf mRNA levels, one of the most prevalent neurotrophic factors  
25 in the brain. This increase in Bdnf is in contrast to reports of acute and chronic stress decreasing levels of Bdnf in hippocampus (11).

- In conclusion gene expression profiling of several brain areas of CRH overexpressing mice elucidated previously unrecognized homeostatic mechanisms present in these  
30 animals. The identification of genes altered in CRH overexpressing mice compared to wild-type littermates allow further molecular characterization of this animal model of chronic pituitary-adrenal activation. We provide the tools to evaluate at the level of gene expression the consequences of prolonged exposure to CRH, which could eventually lead to the use of novel markers in stress, anxiety or depression both in animals and in humans. In addition we demonstrate that administration of a CRH-R1  
35 antagonist does not lead to a general 'normalization' of expression levels in transgenic animals, although at the pituitary level some effect of the compound is measured. In addition repeated stress inflicted by the treatment seems to elicit a response in gene

-47-

expression of specific genes such as neurotensin receptor 1. This effect can be reversed by CRH-R1 specific antagonists, suggesting a close link between neurotensin and CRH.

## REFERENCES

- 5 1. De Souza EB 1995 Corticotropin-releasing factor receptors: physiology, pharmacology, biochemistry and role in central nervous system and immune disorders. *Psychoneuroendocrinology* 20:789-819
2. Holsboer F 2001 Stress, hypercortisolism and corticosteroid receptors in depression: implications for therapy. *J Affect Disord* 62:77-91
- 10 3. Holsboer F, Gerken A, Stalla GK, Muller OA 1987 Blunted aldosterone and ACTH release after human CRH administration in depressed patients. *Am J Psychiatry* 144:229-231
4. Holsboer F, Gerken A, von Bardeleben U, Grimm W, Beyer H, Muller OA, Stalla GK 1986 Human corticotropin-releasing hormone in depression--correlation with thyrotropin secretion following thyrotropin-releasing hormone. *Biol Psychiatry* 21:601-611
- 15 5. Nemeroff CB, Owens MJ, Bissette G, Andorn AC, Stanley M 1988 Reduced corticotropin releasing factor binding sites in the frontal cortex of suicide victims. *Arch Gen Psychiatry* 45:577-579
6. Raadsheer FC, Hoogendijk WJ, Stam FC, Tilders FJ, Swaab DF 1994 Increased numbers of corticotropin-releasing hormone expressing neurons in the hypothalamic paraventricular nucleus of depressed patients. *Neuroendocrinology* 60:436-444
- 20 7. Wouters L, Göhlmann HW, Bijnens L, Kass SU, Molenberghs G, Lewi PJ 2002 Graphical exploration of gene expression data: a comparative study of three multivariate methods. *Biometrics*
8. Tusher VG, Tibshirani R, Chu G 2001 Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* 98:5116-5121
- 25 9. Van Pett K, Viau V, Bittencourt JC, Chan RK, Li HY, Arias C, Prins GS, Perrin M, Vale W, Sawchenko PE 2000 Distribution of mRNAs encoding CRF receptors in brain and pituitary of rat and mouse. *J Comp Neurol* 428:191-212
- 30 10. Tsai KJ, Chen SK, Ma YL, Hsu WL, Lee EH 2002 *sgk*, a primary glucocorticoid-induced gene, facilitates memory consolidation of spatial learning in rats. *Proc Natl Acad Sci U S A* 99:3990-3995

-48-

11. Heinrichs SC, Stenzel-Poore MP, Gold LH, Battenberg E, Bloom FE, Koob GF, Vale WW, Pich EM 1996 Learning impairment in transgenic mice with central overexpression of corticotropin-releasing factor. Neuroscience 74:303-311

5

SEO ID NO.1

SEQ ID No.2

15      **SEO ID No. 3**

20      SEO ID No.4

25

## SEO ID No.5

SEO ID No.6

40      SEO ID No.7

SEQ ID No.8

55      SEO ID No.9



55

55

SEO ID No.12

15

20 cccagagatagaagctgacgctggcgccaaacccctgggggaaaggttgctcgggcaagtagtcatggctgaaagcagctgggaatcgtataaagacaaacc  
agggagggcgggtcaccgtggcagtggaagatgttgaagatgaatgccaaagaggaagggaccgtctgactctggatcagagatggngatgatgaagatgattg  
ggaaacatagaacatattacaacccctcctggggggcctgcacgcaaggatggaccctctctacgtcatalagtgatattgcatcgonnaggcaacccctcggggaata  
cctccgagcccggaaggccacctggcatggaggaactcctatgacattaacctgtgtcccggaggagcagatgacctcaagggacttgggtgtcctgcaccta  
ccagctggcttagaggcagtggaagtacttggcttcccaaaaatgtatccatcgagatattggcctgcagaaacgtgttggttaacgannacaatgtgatgaag  
25 atagcagagcttggcctggccaggggatacaacaacatgactactataaaagaccacaaatgtggcgacgtccagctcaagtggtatggctcctggaagcc  
cttttgatagagtttacacatcagagggatgtctgtgcttcgggggtgttaatgtgggagatcttacttttgggggtccaccttaccagggattcccg  
ggaggaaacttttaagctgctcaaaaggaggacacaggatggacaaggcccaaccaactgcaccnaigaaactgtatacgtatgatagggaattgctggcatgc  
tgtaacctcacagagaccacattcaagcaggttggctgaagacttgatcgaattcgaactcaaacacnattaggggaattacttggatctcaccagccctc  
cgaacagagtattctctagttaccccgaacaagatgctctgtgttcacgggagcagattctgtgttttccagaccccaatgaccttatgaacccctgtcgtcca  
rtatccacacataaagcgacagtttataacatga

35 PRDKLTLGKPLGEGCFGQVVM AEAVGIDKDKPKEAVTVAVKMLKDDATEKDLSDLVSEMEM  
MKMIGKHKHNINLLGACTQDGPLYVIVEYASKGNLREYLRARRPPGMEYSYDINRVPEEQMTFK  
DLVSCYQLARGMEYLASQKCIHRDLAARNVLVTENNVMKIADFLGLARDINNIDYKKTNGR  
LPVKWMAPEALFDRVYTHQSDVWSFGVLMWEIFTLGGSPYPGPVVEELFKLLKEGHRMDKPTN  
CTNELVMMMRDCWHVAPSQRPTFKQLVEDLDRILTLTTNEEYLDLTQPLEQYSPSYPTSSCS  
SGDSDVFSPPDMPYEPCLPQYPHINGSVKT

[illegible]

-53-

gcttcagggtccagcccaagaccccccagggagggccaggttcctgtgtgactgtgaaccacccccgggtggatnaggatctcaagccagacagaaaa  
 gtcaggccggcacccttagacataaaacctctgtgaatgggnaagagctgcaagcccccagtcagatctccatcaocaggagcttcgctcgggactcc  
 tctaggttcccatgtccctcggcctgattctgtgcacagtagacatcggagcagcagctcctatgacagtgaaagaaactatgtcccatgaatccaaat  
 ctgtctgagcgaagaccgaatctctgtccagcaacagccttgaatgggnaagcagcccgatgaatanaacccaagggagacaaaacagctgaatacct  
 5 gggattagacactgttctgggaagtcacgccaccacgganagcaaaagagcaggtgtgtcggcagcagcatggcagacgaggggtggattacgtg  
 tgggtggaccacagaaagactctggccctgaagagtaccagagaagccttggacggatgggagggcagtcacagagtcggagacacccacaagaatg  
 tgaagtgaagacatgccgtcgcctctgagggaagacgagatctgagtggaagagagatgccaaagaaagatgttccactctcagtgaggcctcggag  
 cagagggggaagaggaaggaatcctcacacatgttcaagcaaaatgggttgaatgtgtcgtgtgtgtattgattataacgtgtanaaaacccgggga  
 aatagtgttttagttcacagagaagccttctgcccataaacaacacctgtatgattataaacaactaagtgaaactgagctactgttaattcggaaattcaggt  
 10 atciaccttaacagacttccctgggagggtctttggcctccacacactactatataaacaactaagtgaaactgagctactgttaattcggaaattcaggt  
 gaagctacaggggctaacaccattaaacaagaagtgnaagtgacacattcgcctttcttgaaggtggtagccattagcttaagctgtgaacatagttggc  
 cg

SEQ ID No.16

MSGGEVVC SGWLRKSPPEKKLKRYAWKRRWFVLRSGRLTGDPDVLEYYKNDHAKKPIRIIDLN  
 LCQQVDAQLTFNKKEFENSYIFDINTIDRIFYL VADSEEDMNK WVRICDICGFNPTEEDPVKPLT  
 QSSQAPVDSPPFAISTAPASSQMEASSVALPPPYQVISLPPHPDTLGLQDDPDQYLLINCQSKKPEP  
 20 NRTLFD SAKPTFSETDCNDNVPSHQTPASSQSKHGMNGFLQQMMYDCPPSRLTSVSGESSLYN  
 LPRSYSHDVLPKESPSSTEADGELYTFNTPSGTAGVETQMRHVSISYDIPPTPGNTYQIFRTFFEST  
 LGQSSKLDITFDIPPRPPKPHPSHDRSPVETCOVPTASDTSYCIFFPVGMTFPRSNTISTVDLN  
 KLRKDASSQDCYDIPRTFFPSDRSSSLEGFHSQYKIKSVLTAGGVSGEELDENVFPMNPNSPFRQH  
 SSGSFYQEPNYVPMTPGTDFSSFGMQVPPPAHMOFRSSPKTFFRRPVVADCEPPPVDNRNLPK  
 25 DRKVKPAFLDIKPLSEWEELOAPVRSPTIRSFA RDSRRFPMSPRPDSVHSTSSSDSHDSEENYVP  
 MNPNLSGEDPNLFASNSLDGSSPMNPKPKGDKQVEYLDL DLDLSGKSTPPRKQKSSGSGSSMADE  
 RVDYVVVDQQKTLALKSTREAWTDGRQSTESETPTKNVK

SEQ ID No.17

gtcgaaccacgcgtccggggcggcggtcgggtgtgtggacctgacagcggccgtcggccacacctogctctctcctggccggcggtatcctttgtct  
 ctcagagcggcggaaggtcgcagagcagcgcggcggtcggggacgcaaacgcccgtctctcgggtcactcccccagcctcggccaccgcagaaac  
 ttgggctcagagaagccagcggacccgcgtcagctcagagcgcgggtttagtctccactgccaagcgtccatgggcaagcagaaacagcgtcgtcct  
 35 gnggtgtcagagcagcgtcgggaacacacgggaattcaccgacccatgaagcttcagaggtgtgtacnngggcttctcaaggactcggccacgtgcccacot  
 gactgtggacgagttcaagaagatctacgccaacttctccctacggcgatgcctcgaagttcgtgaacacgcttccgcaccttcgacacccaacagc  
 gatggcaccatcgaatcgggaggttcatcattgtctgagtgtagcttcgggggaagcgtggaacagaagctcaagtgggcccttagcatgtacgacct  
 ggacgggaattggctacatcagccgcaaggaatgtcggagatagtcagggccntctacaagatggtgtcctccgtgatgaagatgcttgaggacgagt  
 ccacggccggagagcgaacagaaagatctcnggcagatggacacaaacnaatgatggcaaaactgtccctgggaagaattcatnaaagggtgcaagag  
 40 cgacccatccattgtccgggtgtcgtcagtgtagtccagcagtgctagccaggtcgtgaggggaatgggcttggacatttggagaaacacagggcttggc  
 ctggccattctcagcttgcgtgnaaagtggaattctcgtgactctctcgtctcccccaggaactgggtccaggcagcaatcgacccctcctgacctggcca  
 actgtggcttccctcctcctcagtcgaaggtcggctcctcccaatggttcnaagtggtcctcgggtgtcgggtcctcctgcccacagagggcctggcgccct  
 taaggacaccagacaggaactttagtatcttaccagaccagcctaattatctatgacccaggtgactttgagggatgaaaaagatggagagtagaca  
 45 ggcaagccctcctgtgcatggctccacctccctgttcatcttctcgaacaaagccgttgcattatataaccgtggatccttactttaalatatagattat  
 gtatggtgagatggactaataatgtcagccctgtatitaaatgctcgtgactgcttgaagcagtggtccccgtggcctgtgattccccccaccccaagct  
 ggctgtgtgtgtgtatgtatgattcatgattgcttaccacnngagatcatgcatgccagggcgtgtggacatctgcccctcgtctgnaatgacgg  
 ttttctgtgtgtcctcctcgtcgtattggtatggcatattaaatgatcaaat

SEQ ID No.18

MGKQNSKL RPEVLQDLREHTEFTDHELQEWYKGF LKDCPTGHLTVDEFKKIYANFFPYGDASK  
 FAEHVFRFTD TNSDGTIDFREFTIALSVTSRGKLEQKLK WAFSMYDL DNGYISRSEM LEIVQATY  
 KMVSSVMKMP EDESTPEKRTDKIFRQMDT NNDGKLSLEEFK OAKSDPSIVRLQCDPSSASQF

SEQ ID No.19

55      SEQ ID No.23

[illegible]

10 · SEQ ID No. 24

MEKTELIQKAKLAEQAERYDDMATCMKAVTEQGAELSNEERNLLSVAYKNVVGGRRSAWRVI  
SSIEQKTDTSDDKLQLIKDYREKVESELRSICTTVLELLDKYLIANATNPESKVFYLMKGDYFR  
YLAEVACGDDRKQTIENSQGA YQEA FDISKKEMQPTHPIRLGLALNFSVFYYEILNNPELACTLA  
KTA FDBAIAELDTLNEDSYKDSTLIMQLLRDNLTLWTSDSAGEECDAAEGAEN

SEQ ID No.25

20 **ttttttttttttcccatcccttgctggccttatctggggcagggggaaacgatctctggcagtggtaggacatggaggtacaaagagttggaaagccctgggg**  
**gctcagctttcacaggcaggggtgggcacacttgcgaagtgccagagagggagggccgttatatgtcaacccacaggagaaatatctctatgcagccgtt**  
**cattatgctcatgtagcagggacaggtgtgcgggtcatgtggcgagaaagcccaagagcttgggataccccaccgcccggaataagccggctgtcagtggcagc**  
**agggaaaagatctotgctccagagtgagggttcatttgcccttgcaaggctccgcgtgaaggtgccaccaggagatctgaatcatccgtgggtgtcattcagaggc**  
**tctgtaaactcttga**

**SEQ ID No.26**

[illegible]

**SEQ ID No.27**

gtgacaggacgcccacaccccatcttaagctttatctctagtagcctgaggggctggagagaggtagttctgccggattgtgtcactggagtcgt  
tccgtgggctcctttttctcgtgttgcaattttcttggtccgatgaanagcaattcttttctaaccataaangtgcctcttcagcaatgg

SEQ ID No.28

45 agcaatgaatgttggaacatgaagttaaacctctggtggagggaaattcatcgccctgggttcacagaaatgctgatgggaaaattaaagtgtgaagtttggggctct  
cttcagggatgacagatgtgccaatctctttgaaagcgtttggtaggaacctctgaaagctgcgaaaacgaagggaagattgtacatcagcagggggaactacttt  
gcaaagtggttcatgntgatgttgacattgatttgcgcaggaattaaatgtgggttgacagctctgggtgtatcgtgataactgggaataactaaagttaagagacatgac  
gtgaattctcatgattttatagaacttggattgaacaaaggggggctgttgagagaagctctgtttatataccttgaugcaaaaacattacnaatgtaaaatgagac  
aaacctatatttttcttaagaaaggtaatttgggaaatgtgagttatgaannacttttgggagagtggttggaanaaac

SEQ ID No. 29

attgggcatgtgtgtgctagcaccctgtccatgagaatatttatgatattaaagaaacctttgaaatggcctgtcttcnaaggaagataattcattggcttc  
cattttcagtcctcttcnaaaggctggcctggcctgttttaagactgcagagttgttatgttttaatttgaattatggcctattaaagataggacaaacttggagatt  
atgatgtgtcctggcacagactgtattaaaacaacactcccgtga

55      SEO ID No.30





TGAAGCTGGCGGGGTGCTGACCGGGATGAAAGTGAAGGCCGCAACAGGGGGCGCAACGGTCCATTTTAAATGAGGGAGGAGCTTGAAGAAACAGGGCCAAACGCTATG  
 GCGGGCTTTTGGCCAAATACCCCAAGAGGAGTTCGGAGATGGCCCGGGGATGAGGACGGGGGGCCAGGATGGGGGATCAGGTAAGGGCATGAGGACCTGT  
 ACMAACGCTATGGGGGGCTTCCTGCGGGCATTCGCCCCAAGCTGAAGTGGGAGCAACCAAGAGCGCTATGGTGTGTTCTTCGCGGCGTCAAGTTCAGGCTGG  
 TGACCGGGTCCGAGGAGAACCCCAATACCTATCTGAAGATTAGATGTTTGA

SEQ ID No.35

MAWSRLMLAACLLVMPFSNVMADCLSLCSLCAVRIQDGPFPINPLICSLEQCQDLVPPSEEWETCR  
GFSSFLTTLTVSGLRGKDDLEDEVALEEGISAHAKLLEPVLKELEKSRLTTSVPBEKFRGLSSSFQN  
GKSELAGADRMNDEAAQGRTVHFNEEDLRKQAKRYGGFLRKYPKRSSEMARDEDGGQDGD  
QVGHELDLYKRYGGFLRRIRPKLKWDNQKRYGGFLRRQFKVVTRSQENPNTYSEDLDV

**SEQ ID No.36**

[illegible]

SEQ ID No.37

MARFLRLCTWLLALGSCILLATVQAECSDQCAKCTYRLVVRPGDINFLACTLECEGQLPSFKIWET  
CKDILLQVSRPEFFWDNIDMYKDSSKQDESHLLAKKYGGFMKRYGGFMKKMDELYPMEPEEEA  
NGGEILAKRYGGFMKKDADEGDTLANSSDLLKELLGTGDNRAKDSHQQUESTNNDNEDMSSKRY  
GGFMRLSLKRSQPLEDEAKELQKRYGGFMRRVGRPEWWMMDYQKRYGGFLKRFABSLPSDEEGE  
NYSKEVPIEKSPRYGGFMRF

SEQ ID No.38

accacgpglccggccgggttcacgtcccccagtcicctttgggctcttccgggcatcgggacgatgaccgtcaaaagccgaggctgctcgaagcac  
ccttacctaciccagaatgaggggaatggtagcgattctcctcgtctttatgaacacagaganggatgggacctgaacgattattacagaaatggccagcaa  
caccatgcatgcnaaacacgctgaaggtcagtcattttgaaaatgtcccaicccagggagccggagccttatgaaacgtctaacccctccctccgccaagtc  
cctctcaacaantcaacctgggtccgctcctcaaacccicacgccnaaacctccgacttcaactttgaaagtgaicgggaaagggcagtttggaagaggttc  
ttctggctaaggcacaaggcagaagaagtattctatgcagctcaaaagtttacagaagaaagccnctcctgaaagaaagaaaggaagcatattatgtcaga  
gggaatgttctgttgaaagaaigtgaagcaccctttcctgggtgggcttcacattcattccagaccgctgacanaacctctacttttcttggaactaattaag  
gttcgagaagctgtcttaccatctccagaggagcgcgtcttctcctgaaaccacggggctcgtactacggagctgaaatagccagtgcccttgggctacatgc  
ctcccaanaacatcgtttatagaagcattaaacctgagaatatctctctagacaccaggggcacatcgtccctactgaccttgggctctgcanaagagaatait  
gagcataacgggacacacatcactctgtgtgtggcagcgcctgacattatgtcctctgaagctcccatagacuaegccgtatgaccggacggtgactgtgtgt  
gtcttggggctgtctgtatgagatgctctacggcttcccccgtttatagccggaaacacggctgagatgacgacaaatattctgaacaagccctccag  
tgaaccanaatatncaancctggcnaaggcacctcclggaaggccctcctgcagaaagaaagccacgaagccctgggtgctcnaaggagacttatggag  
aataagagtcataattcttcttitaatnaactgggaigatctcatnaaagaagattacacccccattaaaccnaaagtggaggtccaggtgacacctcgg  
cactttgatcccgagtattaccgaggagccgggtcccaagctccatcggcagggtccctgacagcatcctgtacggccggtgtgaaaggagcagcagaa  
ggcttctcctggcttctctatgacacctcctgtgtgattcttctctgagtgctcccgggagtggttctgaaggagacttctcagcgttctcaagtggttctctac  
cctttgggtgaggttgccagctgacagaacatttanaaagaatttgcncacctgggaagcttggcagctcgcctgccggcgltggcgcgacgcagcgcg



[illegible]

-59-

5 caagccaggcagcccagcccaggactctacccctgggtttctagaccttaagccattctctgagcccttgatttcctcatctcggcgggaccgggactgtctt  
gaaggactcagcctggaggtacacagaagtgggacacctgaggctgggttagagccnatgccctcttacatgattcctcagaaatgtggtggccattata  
gggggggtccaggcagtcacagaaagtctctgctgtagtctatagcatcttgctgtgtgcttggtgtaattagggacatacatccargcaccaccagacac  
acctgtgaccccgccgggtatcctatgctctgggccaagtgaagaagaataagttctggagggtgtcctgagcctcctggcraggtctcatgagccca  
gtcgtgatgtggaaacaagtctgtgtgcccacaaatactgtcagaatgcccacccctccacactcaataaaaagaacacccctccatattgtctcaataaa  
gaataaagggaagctgtgtatctg

SEQ ID No.41

10 MHLNSSVQQGAPSEPGAQPFHPQFGLETMLLALSLNNGSGNSSESILEPNSNLDVNTDIYSKVLV  
TAVYLALFVVGTVGNSVTAFTLARKKSLQSLQSTVHYHLGSLALSDLLILLAMPVELYNFTWV  
HHPWAFGDAGCRGYFLRDACTYATALNVASLSVERYLAICHPFKAKTLMRSRRTKKFISAIWL  
ASALLAVPMLFTMGLQNRSADGQHPGGLVCTPTVDATVKVVIQVNTFMSFLFPMILISILNTVI  
15 ANKLTVMVHQAAEQGRGVCTVGTHNSLEHSTFNMSIEPGRVQALRHGVLVLRAVVIAFVVCW  
LPYHVRRLMFCYISDEQWTTFLDFYHYFYMLTNALFYVSSAINPILYNLVSANFRQVFLSTLAC  
LCPGWRRRRKKRPTFSRKPNMSNNHAFSTSATRETTY

## CLAIMS:

- 5 1. A method of diagnosing a CRH induced gene expression profile in an individual said method comprising;
- a) obtaining a biological sample of said individual; and
- b) determine the level of gene transcription of a gene comprising a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40.
- 10 2. A method according to claim 1 wherein the biological sample is a bodily fluid or tissue sample
- 15 3. A method according to claims 1 or 2 wherein the level of gene transcription is being assessed using a probe which binds to a polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID NO.5, SEQ ID NO.6, SEQ ID No.7, SEQ ID No.8, SEQ ID NO.9, SEQ ID 11, SEQ ID NO.13, SEQ ID NO.15, SEQ ID NO.17, SEQ ID NO.19, SEQ ID NO.21, SEQ ID NO.23, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30, SEQ ID No.31, SEQ ID NO.32, SEQ ID NO.34, SEQ ID NO.36, SEQ ID No.38 and SEQ ID No.40.
- 20 4. A method according to claim 3 wherein the level of gene expression is determined using microarray technology.
- 25 5. A method of diagnosing a CRH induced gene expression profile in an individual said method comprising;
- a) obtaining a biological sample of said individual; and
- 30

b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said biological sample;

whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

6. A method according to claim 5 wherein the biological sample is a bodily fluid or tissue sample

7. A method according to claims 5 or 6 wherein wherein the amount of protein that modulates CRH signaling is being determined using an antibody which binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

8. A method according to claims 5 or 6 wherein the amount of protein that modulates CRH signaling is being determined by assessing the level of gene transcription of a gene encoding an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

9. A method according to claim 8 wherein the level of gene transcription is being assessed using a probe which binds to a polynucleotide encoding an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID NO.12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

10. A method according to claim 3 wherein the level of gene expression is determined using microarray technology.

11. A method for identifying a compound capable to alter the CRH signalling response in a cell, said method comprising;

- a) contacting said cell with CRH in the presence and absence of said compound;  
b) determine the amount of at least one protein that modulates corticotropin releasing hormone (CRH) signaling in said cell; and  
c) compare the amount of said protein in the presence and absence of said compound;

whereby the protein that modulates corticotropin releasing hormone (CRH) signaling is being selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

12. A method according to claim 11 wherein the cell is a eukaryotic cell such as the murine pituitary corticotroph-derived adenoma cell line AtT-20.

13. A method according to claims 11 or 12 wherein the amount of protein that modulates CRH signaling is being determined using an antibody which binds to a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

14. A method according to claims 11 or 12 wherein the amount of protein that modulates CRH signaling is being determined by assessing the level of gene transcription of a gene encoding an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

15. A method according to claim 14 wherein the level of gene transcription is being assessed using a probe which binds to a polynucleotide encoding an amino acid

sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

5

16. A method according to claims 14 or 15 wherein the level of gene expression is analysed using microarray technology.

10

17. A method for identifying a compound capable to alter the CRH signalling response activity in a cell, said method comprising;

15

a) contacting a cell which expresses at least one protein comprising an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41, with said test compound; and

b) compare the CRH response activity of said cell in the presence and absence of said compound.

20

18. A method according to claim 17 wherein the cell is a host cell capable to express at least one protein having an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

25

19. A method according to claim 18 wherein the host cell is transfected with at least one vector comprising a regulatory sequence.

30

20. A method according to claim 18 wherein the host cell is transfected with at least one vector comprising a polynucleotide sequence encoding an amino acid sequence selected from the group consisting of SEQ ID NO.10, SEQ ID 12, SEQ ID NO.14, SEQ ID NO.16, SEQ ID NO.18, SEQ ID NO.20, SEQ ID NO.22, SEQ ID NO.24, SEQ ID NO.33, SEQ ID NO.35, SEQ ID NO.37, SEQ ID No.39 and SEQ ID No.41.

21. An isolated polynucleotide comprising a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID No.5, SEQ ID No.6, SEQ ID No.7, SEQ ID No.8, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30 and SEQ ID No.31, for use as a marker of CRH signaling in a cell.
22. An isolated polynucleotide according to claim 21 wherein said polynucleotide is mRNA, DNA or cDNA.
23. An isolated polynucleotide consisting of a nucleic acid sequence selected from the group consisting of SEQ ID No.1, SEQ ID No.2, SEQ ID No.3, SEQ ID No.4, SEQ ID No.5, SEQ ID No.6, SEQ ID No.7, SEQ ID No.8, SEQ ID No.25, SEQ ID No.26, SEQ ID No.27, SEQ ID No.28, SEQ ID No.29, SEQ ID No.30 and SEQ ID No.31, for use as a marker of CRH signaling in a cell.
24. A vector comprising an isolated polynucleotide according to any one of claims 21 to 23.
25. A vector according to claim 24 wherein the polynucleotide is operably linked to an expression control sequence.
26. A host cell transfected with a vector according to claims 24 or 25.

## ABSTRACT

## CRH RESPONSIVE GENES IN CNS

---

The present invention relates generally to therapy and diagnosis of depression. In particular this invention relates to the polypeptides as well as to the polynucleotides encoding these polypeptides, wherein said polypeptides are shown to play a central role in mediating the endocrine response to corticotropin releasing hormone. These polypeptides and polynucleotides are useful in the diagnosis, treatment and/or prevention of depression.




-1/11-

Table 1

Gene Name	CDS	SEQ ID	NA	AA	Title	Sequence Derived from
down 170015E05RK			1		RIKEN cDNA 170015E05 gene	AB4237
down 230002F08RK			2		RIKEN cDNA 230002F08 gene	AB123584
down 261000B10ARK			3		RIKEN cDNA 261000B10A gene	AB46337
down 2610371G22RK			4		RIKEN cDNA 2610371G02 gene	AB221684
down 483143D1RK			5		RIKEN cDNA 483143D1 gene	AB28842
down A414D47			6		expressed sequence A414D47	AB123849
down Gene4-47834			7		ESTs. Moderately similar to VIS3 MOUSE VISININ LIKE PROTEIN 3 [M.musculus]	CB0148
down			8		ESTs, moderately similar to VIS3 MOUSE VISININ LIKE PROTEIN 3 [M.musculus]	AB43383
down B-FABP			9		Brain fatty acid-binding protein gene	U04827
down B-FABP			10		Brain fatty acid-binding protein	AA461804
down Ebf2	214-1308		11	12	endothelial differentiation, lysophospholipid acid G-protein-coupled receptor, 2	U13370
down Frl2	1-1038		13	14	fibroblast growth factor receptor, 2	AF223822
down Gdn1	24-2111		15	16	growth factor receptor bound protein 2-associated protein 1	AD48226
down Hpsd1	261-942		17	18	hypocapnia-like 1	AF085182
down Hs611b1	128-1008		19	20	hydroxyesteroid 11-beta dehydrogenase 1	AF3202
down Nls2	27-1280		21	22	neurotensin receptor 2	U51938
down Nlsr1	444-1718		40	41	neurotensin receptor 1	NM_01870
down Yohas1	24-758		23	24	tyrosine 3-monooxygenase/tyrosophan G-monooxygenase activation protein, theta polypeptide	U58243
up 111004A122RK			25		RIKEN cDNA 111004A122 gene	AB52940
up 120003D14RK			26		RIKEN cDNA 120003D14 gene	AB218938
up 1600023Q2RK			27		RIKEN cDNA 1600023Q02 gene	AV019397
up 311003A17RK			28		RIKEN cDNA 311003A17 gene	AA839425
up 3110023F10RK			29		RIKEN cDNA 3110023F10 gene	AI185302
up A457657			30		expressed sequence A457657	AB53024
up A450570			31		expressed sequence A450570	AB46545
up Fkyl6	220-1680		32	33	Fkyl6 binding protein 6 (Fkyl6)	U15838
up Pdm	1747		34	35	prolymphoblastin	AV382284
up Perk1	287-1103		36	37	proprotein convertase 1	MF5491
up Sdk	88-1304		38	39	serum albumin precursor regulated kinase	AB104181

-2/11-


Fig. 1



222



Fig.2 continued



**PO1**

Ripras.: 25% • 13%, Claxari = 100%, Cxatc = 60%, Kcm. = p'dm, Scaf = 70%, RUV = m'atn, CVV = m'atn


PC2



-8/11-

Fig.2 continued

## Spectralmap Pituitary




PC1  
 PC2  
 Repres. = 23% + 10%, Clusters = none, Center = double, Mem. = global, Best = use, RV = mean, CW = mean




-9/11-

Fig. 3






-10/11-

Fig. 4



-11/11-

Fig.5



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**